



www.theoryofgroups.ir

International Journal of Group Theory
ISSN (print): 2251-7650, ISSN (on-line): 2251-7669
Vol. x No. x (201x), pp. xx-xx.
© 201x University of Isfahan



www.ui.ac.ir

FINITE GROUPS OF THE SAME TYPE AS SUZUKI GROUPS

SEYED HASSAN ALAVI, ASHRAF DANESHKHAH* AND HOSEIN PARVIZI MOSAED

Communicated by Alireza Abdollahi

ABSTRACT. For a finite group G and a positive integer n , let $G(n)$ be the set of all elements in G such that $x^n = 1$. The groups G and H are said to be of the same (order) type if $|G(n)| = |H(n)|$, for all n . The main aim of this paper is to show that if G is a finite group of the same type as Suzuki groups $Sz(q)$, where $q = 2^{2m+1} \geq 8$, then G is isomorphic to $Sz(q)$. This addresses the well-known J. G. Thompson's problem (1987) for simple groups.

1. Introduction

For a finite group G and a positive integer n , let $G(n)$ consist of all elements x satisfying $x^n = 1$. The *order type* of G is defined to be the function whose value at n is the order of $G(n)$. In 1987, J. G. Thompson [11, Problem 12.37] posed a problem which is related to algebraic number fields:

Is it true that a group is solvable if its type is the same as that of a solvable one?

This problem links to the set $\text{nse}(G)$ of *the number of elements of the same order* in G . Indeed, it turns out that if two groups G and H are of the same type, then $\text{nse}(G) = \text{nse}(H)$ and $|G| = |H|$. Therefore, if a group G has been uniquely determined by its order and $\text{nse}(G)$, then Thompson's problem is true. One may ask this problem for non-solvable groups, in particular, finite simple groups. In this direction, Shao et al [13] studied finite simple groups whose order is divisible by at most four primes. Following this investigation, such problem has been studied for some families of simple groups [1, 2, 3, 13] including Suzuki groups $Sz(q)$ with q prime [10]. In this paper, we generalize the main result in [10] and prove that

MSC(2010): Primary: 20D06; Secondary: 20E32, 20E34.

Keywords: Suzuki group, Thompson's problem, element order.

Received: 08 March 2017, Accepted: 08 July 2017.

*Corresponding author.

Theorem 1.1. *Let G be a group with $nse(G) = nse(Sz(q))$ and $|G| = |Sz(q)|$. Then G is isomorphic to $Sz(q)$.*

As noted above, as an immediate consequence of Theorem 1.1, we have that

Corollary 1.2. *If G is a finite group of the same type as $Sz(q)$, then G is isomorphic to $Sz(q)$.*

In order to prove Theorem 1.1, we use a partition of Suzuki groups $S := Sz(q)$, where $q = 2^{2m+1} \geq 8$, see Lemma 3.2, that is to say, a set of subgroups H_i of S , for $i = 1, \dots, s$, such that each nontrivial element of S belongs to exactly one subgroup H_i . We use this information to determine the set $nse(S)$ in Proposition 3.3. It is also a main tool to show that 2 is an isolated vertex in the prime graph of a group G satisfying hypotheses of Theorem 1.1, see Proposition 4.1. Then we show that G is neither Frobenius, nor 2-Frobenius group. Finally, we obtain a section of G which is isomorphic to S and prove that G is isomorphic to S .

Finally, we give some brief comments on the notation used in this paper. Throughout this article all groups are finite. Our group-theoretic notation is standard, and it is consistent with the notation in [4, 6, 7]. We denote a Sylow p -subgroup of G by G_p . We also use $n_p(G)$ to denote the number of Sylow p -subgroups of G . For a positive integer n , the set of prime divisors of n is denoted by $\pi(n)$, and if G is a finite group, $\pi(G) := \pi(|G|)$, where $|G|$ is the order of G . We denote the set of elements' orders of G by $\omega(G)$ known as *spectrum* of G . The *prime graph* $\Gamma(G)$ of a finite group G is a graph whose vertex set is $\pi(G)$, and two vertices p and q are adjacent if and only if $pq \in \omega(G)$. Assume further that $\Gamma(G)$ has $t(G)$ connected components π_i , for $i = 1, 2, \dots, t(G)$. The positive integers n_i with $\pi(n_i) = \pi_i$ are called order components of G . Clearly, $|G| = n_1 \cdots n_{t(G)}$. In the case where G is of even order, we always assume that $2 \in \pi_1$, and π_1 is said to be the even component of G . In this way, π_i and n_i are called odd components and odd order components of G , respectively. Recall that $nse(G)$ is the set of the number of elements in G with the same order. In other word, $nse(G)$ consists of the numbers $m_i(G)$ of elements of order i in G , for $i \in \omega(G)$. Here, ϕ is the *Euler totient* function.

2. Preliminaries

In this section, we introduce the some known results which will be used in the proof of the main result.

Lemma 2.1. [8, Theorem 9.1.2] *Let G be a finite group, and let n be a positive integer dividing $|G|$. Then n divides $|G(n)|$.*

The proof of the following result is straightforward by Lemma 2.1. Recall that $nse(G) = \{m_i(G) \mid i \in \omega(G)\}$.

Lemma 2.2. *Let G be a finite group. Then for every $i \in \omega(G)$, $\phi(i)$ divides $m_i(G)$, and i divides $\sum_{j|i} m_j(G)$. Moreover, if $i > 2$, then $m_i(G)$ is even.*

Lemma 2.3 (Theorem 3 in [15]). *Let G be a finite group of order n . Then the number of elements whose orders are multiples of t is either zero, or a multiple of the greatest divisor of n that is prime to t .*

In what follows, recall that $t(G)$ is the number of connected components of the prime graph $\Gamma(G)$.

Lemma 2.4. [5, Theorem 1] *Let G be a Frobenius group of even order with kernel K and complement H . Then $t(G) = 2$, $\pi(H)$ and $\pi(K)$ are vertex sets of the connected components of $\Gamma(G)$.*

A group G is called 2-Frobenius if there exists a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that G/H and K are Frobenius groups with kernel K/H and H respectively.

Lemma 2.5. [5, Theorem 2] *Let G be a 2-Frobenius group of even order. Then $t(G) = 2$, $\pi(G/K) \cup \pi(H) = \pi_1$, $\pi(K/H) = \pi_2$, and G/K and K/H are cyclic groups and $|G/K|$ divides $|\text{Aut}(K/H)|$.*

3. Elements of the same order in Suzuki groups

In this section, we determine the set of the number of elements of the same order in Suzuki groups.

Lemma 3.1 ([14]). *Let $S = \text{Sz}(q)$ with $q = 2^{2m+1} \geq 8$. Then $\omega(S)$ consists of all factors of 4 , $q - 1$ and $q \pm \sqrt{2q} + 1$.*

Let G be a group, and let H_1, \dots, H_t be subgroups of G . Then the set $\{H_1, \dots, H_t\}$ forms a partition of G if each non-trivial element of G belongs to exactly one subgroup H_i of G . Lemma 3.2 below introduces a partition of Suzuki groups.

Lemma 3.2. *Let $S = \text{Sz}(q)$ with $q = 2^{2m+1} \geq 8$, and let $\mathbb{F} := \text{GF}(q)$. Then*

- (a) S possesses cyclic subgroups U_1 and U_2 of orders $q + \sqrt{2q} + 1$ and $q - \sqrt{2q} + 1$, respectively;
- (b) if $1 \neq u \in U_i$, for $i = 1, 2$, then $S(u) = U_i$. Moreover, $|\mathbf{N}_S(U_i) : U_i| = 4$;
- (c) S possesses a cyclic subgroup V of order $q - 1$ and $|\mathbf{N}_S(V) : V| = 2$;
- (d) S possesses a 2-subgroup W of orders q^2 and exponent 4 and $|S : \mathbf{N}_S(W)| = q^2 + 1$. Moreover, the elements of W are of the form

$$(3.1) \quad w(a, b) := \begin{pmatrix} 1 & 0 & 0 & 0 \\ a & 1 & 0 & 0 \\ b & a\pi & 1 & 0 \\ a^2(a\pi) + ab + b\pi & a(a\pi) + b & a & 1 \end{pmatrix},$$

where $a, b \in \mathbb{F}$ and $\pi \in \text{Aut}(\mathbb{F})$ maps x to $x^{2^{m+1}}$, for all $x \in \mathbb{F}$.

- (e) the conjugates of U_1, U_2, V and W form a partition of S .

Proof. All parts of this result follow from Lemma 3.1 and Theorem 3.10 in [9] except for the facts that $|\mathbf{N}_S(V) : V| = 2$ and $|S : \mathbf{N}_S(W)| = q^2 + 1$ which can be found in the proof of Theorem 3.10 in [9]. □

Proposition 3.3. *Let $S = \text{Sz}(q)$ with $q = 2^{2m+1} \geq 8$. Then the set $nse(S)$ consists of exactly one the following numbers*

- (a) $1, (q - 1)(q^2 + 1), q(q - 1)(q^2 + 1)$;
- (b) $\phi(i)q^2(q \mp \sqrt{2q} + 1)(q - 1)/4$, where $i > 1$ divides $q \pm \sqrt{2q} + 1$;

(c) $\phi(i)q^2(q^2 + 1)/2$, where $i > 1$ divides $q - 1$.

Proof. Suppose that $i \in \omega(S)$ is an even number. Then by Lemma 3.1, we have that $i = 2$ or $i = 4$ and i divides the order of subgroup W as in Lemma 3.2(d). Then each element of W is of the form $w(a, b)$ as in (3.1). Obviously,

$$w(a, b)w(c, d) = w(a + c, b + d + (a\pi)c).$$

This in particular shows that $w(0, b)$ (with $b \neq 0$) are the only elements of W of order 2. Therefore, the number of involutions in W is $q - 1$. Since W is a part of the partition introduced in Lemma 3.2(e), the elements of order 2 of S belong to exactly one of the conjugates of W . Thus by Lemma 3.2(d), there are $q^2 + 1$ conjugates of W implying that there are exactly $m_2(S) = (q - 1)(q^2 + 1)$ involutions in S . It also follows from Lemma 3.2(d) that the number of elements of order 4 in W is $q^2 - q$, and hence applying the partition in Lemma 3.2(e), we conclude that S consists of $m_4(S) = q(q - 1)(q^2 + 1)$ elements of order 4. This proves part (a).

Suppose now $i \in \omega(S)$ is an odd number. Then, by Lemma 3.1, i divides the order of one the cyclic subgroups U_1, U_2 and V as in Lemma 3.2, say H . Assume $i = np^\alpha$ with p odd. Since H is a part of the partition introduced in Lemma 3.2(e), the elements of order i are contained in H and its conjugates. Since also H is cyclic, there are $\phi(i)$ elements of order i in each conjugates of H including H .

Now we consider each possibility of H . If $H = U_t$ of order $q \pm \sqrt{2q} + 1$, for $t = 1, 2$, then by Lemma 3.2(b), $|N_S(U_t) : U_t| = 4$, and so $|S : N_S(U_t)| = |S|/4|U_t|$, for $t = 1, 2$. So there are $|S|/4|U_t|$ conjugates of U_t implying that there are exactly $m_i(S) = \phi(i)q^2(q - 1)(q \mp \sqrt{2q} + 1)/4$ elements of order i in S . This follows part (b). If $H = V$, then Lemma 3.2(c) implies that $|N_S(V) : V| = 2$, and so the same argument as in the previous cases, we conclude that $m_i(S) = \phi(i)|S|/2|V| = \phi(i)q^2(q^2 + 1)/2$. This follows (c). \square

4. Proof of the Main Theorem

In this section, we prove Theorem 1.1. From now on, set $S := \text{Sz}(q)$, where $q = 2^{2m+1} \geq 8$, and recall that G is a finite group with $\text{nse}(G) = \text{nse}(S)$ and $|G| = |S|$. Therefore, by Proposition 3.3, $\text{nse}(S)$ consists of

$$\begin{aligned} m_1(S) &= 1; \\ m_2(S) &= (q - 1)(q^2 + 1); \\ (4.1) \quad m_4(S) &= q(q - 1)(q^2 + 1); \\ m_i(S) &= \phi(i)q^2(q \mp \sqrt{2q} + 1)(q - 1)/4, \text{ where } i > 1 \text{ divides } q \pm \sqrt{2q} + 1; \\ m_i(S) &= \phi(i)q^2(q^2 + 1)/2, \text{ where } i > 1 \text{ divides } q - 1. \end{aligned}$$

Proposition 4.1. *The vertex 2 is an isolated vertex in $\Gamma(G)$.*

Proof. Assume the contrary. Then there is an odd prime divisor p of $|G|$ such that $2p \in \omega(G)$. Let $f(n)$ be the number of elements of G whose orders are multiples of n . Then by Lemma 2.3, $f(2)$ is a

multiple of the greatest divisor of $|G|$ that is prime to 2. Since $(q^2 + 1)(q - 1)$ is the greatest divisor of $|G|$ which is coprime to 2, there exists a positive integer r such that $f(2) = (q^2 + 1)(q - 1)r$ and $(r, 2) = 1$. On the other hand, by Lemma 2.2, it is obvious that $m_2(G) = m_2(S)$, and so

$$f(2) = m_2(G) + \sum_{i>2 \text{ is even}} m_i(G),$$

with m_i as in (4.1). Now applying Proposition 3.3, there is a non-negative integer α such that

$$f(2) = (q^2 + 1)(q - 1) + \alpha q(q^2 + 1)(q - 1) + g(2),$$

where

$$g(2) = \sum_{\substack{i|q^2 \pm \sqrt{2q} + 1 \\ i \neq 1}} \beta_i \cdot m_i(S) + \sum_{\substack{i|q-1 \\ i \neq 1}} \gamma_i \cdot m_i(S)$$

for some non-negative integers β_i and γ_i . Since $2p \in \omega(G)$, we have that $\alpha q(q^2 + 1)(q - 1) + g(2) > 0$. Then

$$g(2) = (q^2 + 1)(q - 1)(r - 1 - \alpha q).$$

We now prove that q^2 divides $g(2)$. It follows from Lemma 3.1 that 2 is an isolated vertex of $\Gamma(S)$. Then a Sylow 2-subgroup of S , say S_2 , acts fixed point freely (by conjugation) on the set of elements of order $i \neq 1, 2, 4$ (see Proposition 3.3 and (4.1)). Thus $|S_2|$ divides $m_i(S)$ with $i \neq 1, 2, 4$. Hence q^2 divides $m_i(S)$ implying that $g(2)$ is a multiple of q^2 .

We now consider the following two cases:

(1) Let $g(2) \neq 0$. Then q^2 divides $r - 1 - \alpha q$, and so $q^2 + \alpha q + 1 \leq r$. This implies that $|G| = q^2(q^2 + 1)(q - 1) < (q^2 + 1)(q - 1)r = f(2)$, which is impossible.

(2) Let $g(2) = 0$. Then $r - 1 - \alpha q = 0$ and $\alpha \neq 0$, and so $m_{2p}(G) = q(q^2 + 1)(q - 1)$. Therefore

$$f(p) = \sum_{p|i} m_i(G) = \alpha' q(q^2 + 1)(q - 1) + \sum_{\substack{i|q^2 \pm \sqrt{2q} + 1 \\ i \neq 1}} \beta'_i \cdot m_i(S) + \sum_{\substack{i|q-1 \\ i \neq 1}} \gamma'_i \cdot m_i(S),$$

where α' , β'_i and γ'_i are non-negative integers. Since $m_{2p} = q(q^2 + 1)(q - 1)$, we have that $\alpha' > 0$. On the other hand, by Lemma 2.3, $f(p) = q^2(q^2 + 1)(q - 1)r'/|G_p|$ with r' a positive integer. Thus

$$(4.2) \quad \frac{q^2(q^2 + 1)(q - 1)r'}{|G_p|} = \alpha' q(q^2 + 1)(q - 1) + \sum_{\substack{i|q^2 \pm \sqrt{2q} + 1 \\ i \neq 1}} \beta'_i \cdot m_i(S) + \sum_{\substack{i|q-1 \\ i \neq 1}} \gamma'_i \cdot m_i(S).$$

Since q^2 divides both $q^2(q^2 + 1)(q - 1)r'/|G_p|$ and $m_i(S)$ in (4.2), it follows that q^2 divides $\alpha' q(q^2 + 1)(q - 1)$. Then $q | \alpha'$, and so $|G| = q^2(q^2 + 1)(q - 1) \leq \alpha' q(q^2 + 1)(q - 1) \leq f(p)$, which is impossible. \square

Proposition 4.2. *The group G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that H and G/K are π_1 -groups and K/H is a non-abelian simple group, H is a nilpotent group and $|G/K|$ divides $|\text{Out}(K/H)|$.*

Proof. By Proposition 4.1, the vertex 2 is an isolated vertex in the prime graph $\Gamma(G)$ of G . This implies that the number $t(G)$ of connected components of the prime graph $\Gamma(G)$ is at least two. The assertion follows from [16, Theorem A] provided that G is neither a Frobenius group, nor a 2-Frobenius group.

Let G be a Frobenius group with kernel K and complement H . Then by Lemma 2.4, we must have $t(G) = 2$, $\pi(H)$ and $\pi(K)$ are vertex sets of the connected components of $\Gamma(G)$. By Proposition 4.1, the vertex 2 is an isolated vertex in $\Gamma(G)$. Then either (i) $|K| = q^2$ and $|H| = (q^2 + 1)(q - 1)$, or (ii) $|H| = q^2$ and $|K| = (q^2 + 1)(q - 1)$. Both cases can be ruled out as $|H|$ must divide $|K| - 1$.

Let G be a 2-Frobenius group. Then Lemma 2.5 implies that $t(G) = 2$ and G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that G/H and K are Frobenius groups with kernel K/H and H respectively, $\pi(G/K) \cup \pi(H) = \pi_1$, $\pi(K/H) = \pi_2$ and $|G/K|$ divides $|\text{Aut}(K/H)|$. Since 2 is an isolated vertex of $\Gamma(G)$ by Proposition 4.1, $|K/H| = (q^2 + 1)(q - 1)$ and $|G/K| \cdot |H| = q^2$. Since also K is a Frobenius group with kernel H , there is a positive integer α such that $(q^2 + 1)(q - 1)$ divides $2^\alpha - 1$, which is a contradiction. \square

4.1. Proof of Theorem 1.1.

Proof. Let $S := \text{Sz}(q)$, where $q = 2^{2m+1} \geq 8$. Suppose that G is a finite group with $\text{nse}(G) = \text{nse}(S)$ and $|G| = |S|$. By applying Proposition 4.2, the group G has a normal series $1 \trianglelefteq H \trianglelefteq K \trianglelefteq G$ such that H and G/K are π_1 -groups and K/H is a non-abelian simple group. Since 3 is a prime divisor of all finite non-abelian simple groups except for Suzuki groups. Moreover, 3 is coprime to $|K/H|$. Then $K/H \cong \text{Sz}(q')$, where $q' = 2^{2m'+1}$. This, in particular, implies that $2^{4m'+2}$ divides 2^{4m+2} , and hence $m' \leq m$. On the other hand, H and G/K are π_1 -groups. Then $(q^2 + 1)(q - 1)$ divides $|K/H|$, and so $(q^2 + 1)(q - 1)$ divides $(q'^2 + 1)(q' - 1)$. Since now $m' \leq m$, we must have $m = m'$. Therefore $K/H \cong S$. Now $|G| = |K/H| = |S|$, and hence $G \cong S$. \square

Acknowledgments

The authors are grateful to the anonymous referees and the editor for careful reading of the manuscript and for corrections and suggestions.

REFERENCES

- [1] S. H. Alavi, A. Daneshkhah and H. P. Mosaed, On quantitative structure of small Ree groups, *Communications in Algebra*, **45** no. 9 (2017) 4099–4108.
- [2] A. Khalili Asboei, Characterization of projective general linear groups, *Int. J. Group Theory*, **5** no. 1 (2016) 17–28.
- [3] A. Khalili Asboei and S. S. Salehi Amiri, A new characterization of $PSL(2, 25)$, *Int. J. Group Theory*, **1** no. 3 (2012) 15–19.
- [4] R. W. Carter, *Simple groups of Lie type*, John Wiley & Sons, London-New York-Sydney, **28**, Pure and Applied Mathematics, 1972.
- [5] G. Y. Chen, On structure of Frobenius and 2-Frobenius group, *J. Southwest China Normal Univ.*, **20** no. 5 (1995) 485–487.

- [6] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *Atlas of finite groups*, Oxford University Press, Eynsham, 1985.
- [7] D. Gorenstein, *Finite groups*, Chelsea Publishing Company., New York, second edition, 1980.
- [8] M. Jr. Hall, *The theory of groups*, The Macmillan Co., New York, N.Y., 1959.
- [9] B. Huppert and N. Blackburn, *Finite groups. III*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **243**, Springer-Verlag, Berlin-New York, 1982.
- [10] A. Iranmanesh, H. Parvizi Mosaed and A. Tehranian, *Characterization of suzuki group by nse and order of group*, Bulletin of the Korean Mathematical Society, Accepted, 2015.
- [11] V. D. Mazurov and E. I. Khukhro, *Unsolved Problems in Group Theory, The Kourovka Notebook*, **18** 2014, Available at arxiv.org/abs/1401.0300v6.
- [12] L. Mousavi and B. Taeri, *A characterization of $L_2(81)$ by nse*, Int. J. Group Theory, **5** (1) 29–35 (2016).
- [13] C. Shao, W. Shi and Q. Jiang, *Characterization of simple K_4 -groups.*, Front. Math. China, **3** no. 3 (2008) 355–370.
- [14] W. J. Shi, *A characterization of Suzuki's simple groups*, Proc. Amer. Math. Soc., **114** no. 3 (1992) 589–591.
- [15] L. Weisner, *On the number of elements of a group which have a power in a given conjugate set*, Bull. Amer. Math. Soc., **31** no. 9-10 (1925) 492–496.
- [16] J. S. Williams, *Prime graph components of finite groups*, J. Algebra, **69** no. 2 (1981) 487–513.

Seyed Hassan Alavi

Department of Mathematics, Faculty of Science, Bu-Ali Sina University, Hamedan, Iran

Email: alavi.s.hassan@basu.ac.ir

Email: alavi.s.hassan@gmail.com

Ashraf Daneshkhah

Department of Mathematics, Faculty of Science, Bu-Ali Sina University, Hamedan, Iran

Email: adanesh@basu.ac.ir

Email: daneshkhah.ashraf@gmail.com

Hosein Parvizi Mosaed

Alvand Institute of Higher Education, Hamedan, Iran

Email: h.parvizi.mosaed@gmail.com