



www.combinatorics.ir

Transactions on Combinatorics

ISSN (print): 2251-8657, ISSN (on-line): 2251-8665

Vol. 7 No. 4 (2018), pp. 11-24.

© 2018 University of Isfahan



www.ui.ac.ir

DEGREE RESISTANCE DISTANCE OF TREES WITH SOME GIVEN PARAMETERS

FANGGUO HE AND XINNONG JIANG*

Communicated by Ivan Gutman

ABSTRACT. The degree resistance distance of a graph G is defined as $D_R(G) = \sum_{i < j} (d(v_i) + d(v_j))R(v_i, v_j)$, where $d(v_i)$ is the degree of the vertex v_i , and $R(v_i, v_j)$ is the resistance distance between the vertices v_i and v_j . Here we characterize the extremal graphs with respect to degree resistance distance among trees with given diameter, number of pendent vertices, independence number, covering number, and maximum degree, respectively.

1. Introduction

The graphs considered in this paper are simple and undirected. Let $G = (V(G), E(G))$ be a graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G) = \{e_1, e_2, \dots, e_m\}$. The distance $d(v_i, v_j)$ between the vertices v_i and v_j of the graph G is the length of the shortest path between v_i and v_j . If $S \subseteq V(G)$, let $G - S$ be the subgraph of G obtained by deleting the vertices of S and the edges incident with them. Similarly, for $E \subseteq E(G)$, we denote by $G - E$ the subgraph of G obtained by deleting the edges of E . Let $P_k = v_1v_2 \cdots v_k$ ($k \geq 2$) be a path of G with distinct vertices v_1, v_2, \dots, v_k and assume that $d_G(v_1) \geq 3, d_G(v_2) = \cdots = d_G(v_{k-1}) = 2$, then P_k is called a pendent path of G if $d_G(v_k) = 1$, and P_k is called an internal path of G if $d_G(v_k) \geq 3$. Other undefined notations and terminology from graph theory can be found in [3].

MSC(2010): Primary: 05C15; Secondary: 05C69.

Keywords: Trees, Degree resistance distance, Diameter, Covering number.

Received: 18 December 2017, Accepted: 11 August 2018.

*Corresponding author.

DOI: <http://dx.doi.org/10.22108/toc.2018.108656.1538>

The sum of the distances between all pairs of vertices in a graph G is the Wiener index, namely, $W(G) = \sum_{i < j} d(v_i, v_j)$. It is introduced in 1947 [29] and is one of the most thoroughly studied distance based graph invariants [6, 7, 13, 20, 24]. A modified version of the Wiener index is the degree distance defined as

$$D(G) = \sum_{i < j} (d(v_i) + d(v_j))d(v_i, v_j)$$

where $d(v_i) = d_G(v_i)$ is the degree of the vertex v_i of the graph G . For more information on degree distances of graphs, see [1, 26, 27, 9, 18, 22].

In 1993, Klein and Randić [21] introduced a distance function named resistance distance on a graph. They viewed a graph G as an electrical network such that each edge of G is assumed to be a unit resistor, then take the resistance distance between vertices v_i and v_j to be the effective resistance between them, denoted by $R(v_i, v_j)$. The Kirchhoff index $Kf(G)$ of G is defined as $Kf(G) = \sum_{i < j} R(v_i, v_j)$. The index has been extensively studied in mathematical, physical and chemical aspects (see [2, 15, 31, 12, 33, 34]).

In analogy with the degree distance of a graph, the degree resistance distance (DR-index) of a graph G was first proposed by Gutman, Feng and Yu [14] as

$$D_R(G) = \sum_{i < j} (d(v_i) + d(v_j))R(v_i, v_j)$$

Palacios [25] called this graph invariant as the additive degree-Kirchhoff index. It was systematically studied in [19, 10]. Tu and Su [28] characterized the unicyclic graphs with maximum degree resistance distance, Chen et al. [5] characterized n -vertex unicyclic graphs with given girth having maximum and second maximum DR-index, Yang and Klein gave the formula for the degree resistance distances of the subdivisions and triangulations of graphs [32], In [14] some properties of DR-index were presented and the unicyclic graphs with the minimum and the second minimum DR-index were also completely characterized. The extremal graphs on degree resistance distance of cacti are determined by Liu and Du et al. [23, 10].

If G is a tree, then $R(u, v) = d(u, v)$ for any two vertices u and v [21]. Consequently, the degree distances and degree resistance distances are equal as well [8], i.e.

$$(1.1) \quad D_R(G) = D(G) = 4W(G) - n(n-1) = 4Kf(G) - n(n-1).$$

According to (1.1), if a result holds for DR-index of trees, then it is true for degree distance of trees. Further, the extremal graphs corresponding to the indices $D_R(G)$, $D(G)$, $W(G)$ and $Kf(G)$ of a tree G are the same.

Motivated from [16, 30, 11, 17], we study some mathematical properties of $D_R(G)$ for a graph G . In this paper, we characterize the extremal graphs with respect to degree resistance distance among trees with given diameter, number of pendent vertices, independence number, covering number, and maximum degree, respectively.

2. Preliminaries and some lemmas

Let $R_G(u, v)$ denote the resistance distance between u and v in a graph G . It is known that $R_G(u, v) = R_G(v, u)$ and $R_G(u, v) \geq 0$ with equality if and only if $u = v$.

For a vertex v in G , we define the following two functions

$$Kf_v(G) = \sum_{u \in G} R_G(u, v), \quad D_v(G) = \sum_{u \in G} d_G(u)R_G(u, v).$$

For the sake of conciseness, we write $u \in G$ instead of $u \in V(G)$. By the definition of $D_R(G)$, we also have

$$D_R(G) = \sum_{v \in G} (d_G(v) \sum_{u \in G} R_G(u, v)).$$

Lemma 2.1. [21] *Let G be a graph, x be a cut vertex of G and let u, v be vertices belonging to different components which arise upon deletion of x . Then $R_G(u, v) = R_G(u, x) + R_G(x, v)$.*

Lemma 2.2. [14] *Let G be a connected graph with a cut-vertex v such that G_1 and G_2 are two connected subgraphs of G having v as the only common vertex and $V(G_1) \cup V(G_2) = V(G)$. Let $n_1 = |V(G_1)|, n_2 = |V(G_2)|, m_1 = |E(G_1)|, m_2 = |E(G_2)|$. Then*

$$D_R(G) = D_R(G_1) + D_R(G_2) + 2m_2Kf_v(G_1) + 2m_1Kf_v(G_2) + (n_1 - 1)D_v(G_2) + (n_2 - 1)D_v(G_1).$$

Let v be a vertex of degree $p + 1$ in a graph G , such that vv_1, vv_2, \dots, vv_p are pendent edges incident with v , and u is the neighbor of v distinct from v_1, v_2, \dots, v_p . We form a graph $G' = \sigma(G, v)$ by deleting the edges vv_1, vv_2, \dots, vv_p and adding new edges uv_1, uv_2, \dots, uv_p . We say that G' is a σ -transform of the graph G (see Figure 1).




FIGURE 1. The graphs G and G'

Lemma 2.3. [14] *Let $G' = \sigma(G, v)$ be a σ -transform of the graph G , $d_G(u) \geq 1$. Then $D_R(G) \geq D_R(G')$. Equality holds if and only if G is a star with v as its center.*

Lemma 2.4. *Suppose that G is a connected graph with a cut vertex u and $d_G(u) \geq 3$. Let the paths $P = u_1u_2 \dots u_k$ and $Q = v_1v_2 \dots v_t$ ($k \geq t$) be the connected components of $G - u$, and let $G' = G - v_{t-1}v_t + u_kv_t$ (as shown in Fig.2). Then $D_R(G) < D_R(G')$.*



FIGURE 2. The graphs G and G'

Proof. For the sake of brevity, for any two vertices u, v of G (or G'), we let $d(v) = d_G(v)$ (or $d'(v) = d_{G'}(v)$) and $R(u, v) = R_G(u, v)$ (or $R'(u, v) = R_{G'}(u, v)$). Let $A = \{v_t, v_{t-1}, u_k\}$, $B = \{v_{t-1}, \dots, v_1, u, u_1, \dots, u_{k-1}\}$, $H = V(G) \setminus (A \cup B)$. In the transformation from G to G' , for any pair of vertices x, y satisfying $x, y \in B \cup H$, the term $\sum_{x,y} (d(x) + d(y))R(x, y)$ does not change, and $V(G) = V(G')$. Then

$$\begin{aligned}
 D_1 &= \sum_{x,y \in B \cup H} [(d'(x) + d'(y))R'(x, y) - (d(x) + d(y))R(x, y)] = 0 \\
 D_2 &= \sum_{x,y \in A} [(d'(x) + d'(y))R'(x, y) - (d(x) + d(y))R(x, y)] \\
 &= [2(k + t) + 3 + 3(t - 1 + k)] - [3 + 2(t + k) + 3(k + t - 1)] = 0
 \end{aligned}$$

Note that $R'(u_k, y) = R(u_k, y)$, $R'(v_{t-1}, y) = R(v_{t-1}, y)$ and $R'(v_t, y) = t + k + 1 - R(v_t, y)$ for $y \in B$, let $d_G(u) = d$, then

$$\begin{aligned}
 D_3 &= \sum_{x \in A, y \in B} [(d'(x) + d'(y))R'(x, y) - (d(x) + d(y))R(x, y)] \\
 &= \left(\sum_{x \in A, y \in B - u} + \sum_{x \in A, y = u} \right) [(d'(x) + d'(y))R'(x, y) - (d(x) + d(y))R(x, y)] \\
 &= \sum_{y \in B - u} [4R(u_k, y) + 3R(v_{t-1}, y) + 3(t + k + 1 - R(v_t, y)) \\
 &\quad - (3R(u_k, y) + 4R(v_{t-1}, y) + 3R(v_t, y))] + (2 + d)(k - t + 1) \\
 &= \sum_{y \in B - u} (R(u_k, y) - R(v_{t-1}, y) - 6R(v_t, y)) + 3(t + k + 1)(t + k - 3) + (2 + d)(k - t + 1) \\
 &= 4(t - k - 1) + (2 + d)(k - t + 1) > 0.
 \end{aligned}$$

Since $d'(y) = d(y)$ for $y \in H$, by lemma 2.1, we have

$$\begin{aligned}
 D_4 &= \sum_{x \in A, y \in H} [(d'(x) + d'(y))R'(x, y) - (d(x) + d(y))R(x, y)] \\
 &= \sum_{y \in H} [(1 + d(y))(k + 1 + R(u, y)) + (1 + d(y))(t - 1 + R(u, y)) + (2 + d(y))(k + R(u, y)) \\
 &\quad - (1 + d(y))(t + R(u, y)) - (2 + d(y))(t - 1 + R(u, y)) - (1 + d(y))(k + R(u, y))] \\
 &= \sum_{y \in H} (k - t + 1)(2 + d(y)) > 0.
 \end{aligned}$$

Therefore, by the definition of $D_R(G)$, we get

$$D_R(G') - D_R(G) = D_1 + D_2 + D_3 + D_4 > 0.$$

This completes the proof. □

Let G be a tree, and v be a vertex of G such that $N_G(v) = \{v_1, v_2, \dots, v_{s-1}, u_1\}$. It is obvious that $G - v$ has s ($s \geq 3$) connected components, denoted by T_1, T_2, \dots, T_s . Let there exists a connected component T_i ($i = 1, \dots, s$) such that T_i is a path, without loss of generality, say $T_s = P_k$. Now we construct a new graph G' obtained from G by removing the edges $vv_2, vv_3, \dots, vv_{s-1}$ and adding new edges $v_1v_2, v_1v_3, \dots, v_1v_{s-1}$, as shown in Figure 3. We call the process from G to G' a graph transformation.




FIGURE 3. The graphs G and G'

Lemma 2.5. *Let G and G' be the trees depicted in figure 3. If $d_G(v, u_k) < \max\{d_G(v_1, x) : x \in V(T_1)\}$ or $|V(T_1)| > k + 1$, then $D_R(G') < D_R(G)$.*

Proof. Let $G_1 = G - T_1 - T_s$, $G_2 = G - T_2 - \dots - T_{k-1}$. It is easy to see that that G_1 and G_2 are two connected subgraphs of G and G' having v and v_1 as the only common vertex, respectively. Let $n_1 = |V(G_1)|$, $n_2 = |V(G_2)|$, $m_1 = |E(G_1)|$, $m_2 = |E(G_2)|$. Using Lemma 2.2, we have

$$\begin{aligned} D_R(G) &= D_R(G_1) + D_R(G_2) + 2m_2Kf_v(G_1) + 2m_1Kf_v(G_2) + (n_2 - 1)D_v(G_1) + (n_1 - 1)D_v(G_2) \\ D_R(G') &= D_R(G_1) + D_R(G_2) + 2m_2Kf_{v_1}(G_1) + 2m_1Kf_{v_1}(G_2) + (n_2 - 1)D_{v_1}(G_1) + (n_1 - 1)D_{v_1}(G_2) \end{aligned}$$

By the definition of $Kf_v(G)$ and $D_v(G)$, we know that $Kf_v(G_2) = Kf_{v_1}(G_2)$, $D_v(G_2) = D_{v_1}(G_2)$. Let $H = V(T_1) \setminus \{v_1\}$, $|H| = h$, therefore

$$\begin{aligned} D_R(G) - D_R(G') &= 2m_2(Kf_v(G_1) - Kf_{v_1}(G_1)) + (n_2 - 1)(D_v(G_1) - D_{v_1}(G_1)) \\ &= 2m_2\left[\left(\sum_{x \in H} (R(x, v_1) + 1) + 1 + \frac{k(k+1)}{2}\right) - \left(\sum_{x \in H} R(x, v_1) + \frac{(k+2)(k+1)}{2}\right)\right] \\ &\quad + (n_2 - 1)\left[\left(\sum_{x \in H} d(x)(R(x, v_1) + 1) + d(v_1) + k^2\right) - \left(\sum_{x \in H} d(x)R(x, v_1) + (k+1)^2\right)\right] \\ &= 2m_2(h - k) + (n_2 - 1)\left(\sum_{x \in H} d(x) + d(v_1) - 2k - 1\right) \end{aligned}$$

Note that $\sum_{x \in H} d(x) = 2(h - 1)$, $m_2 = n_2 - 1$, thus

$$D_R(G) - D_R(G') = m_2(4h - 4k + d(v_1) - 3)$$

Since $d_G(v, u_k) < \max\{d_G(v_1, x) : x \in V(T_1)\}$, there exists a vertex $z \in V(T_1)$ such that $d_G(v, u_k) < d_G(v_1, z)$. Thus $h \geq k + 1$. Similarly, if $|V(T_1)| > k + 1$, then $h \geq k + 1$. It follows that

$$D_R(G) - D_R(G') \geq m_2(d(v_1) + 1) > 0.$$

This completes the proof. □

Let G be a connected graph and $P = u_0u_1 \cdots u_{k-1}u_k$ be an internal path of G . Suppose that $G - \{u_1, u_2, \dots, u_{k-1}\} = G_1 \cup G_2$ with $u_0 \in G_1$ and $u_k \in G_2$. We construct a new graph G' obtained from G by contracting the path P and attaching the path P to the common vertex u_0 of G_1 and G_2 , as shown in Fig.4. We call the procedure of constructing G' from G the A-transformation of G .




FIGURE 4. The graphs G and G'

Lemma 2.6. *Assume that G and G' be two connected graphs, and G is an A-transformation of G' . Then $D_R(G) > D_R(G')$ (see Fig.4).*

Proof. Let $A = \{u_0, u_1, \dots, u_{k-1}, u_k\}$, $B = V(G_1) \setminus u_0$ and $C = V(G_2) \setminus uk$. In the transformation from G to G' , the term $\sum_{i,j}(d_G(i) + d_G(j))R_G(i, j)$ does not change for any $i, j \in A$, or $i, j \in B$, or $i, j \in C$. Since $R(i, j) > R'(i, j)$, $d(i) = d'(i)$, $d(j) = d'(j)$ for $i \in B, j \in C$, and $R(i, j) = R'(i, j)$, $d(i) = d'(i)$, $d(j) = d'(j)$ for $i \in A \setminus \{u_0, u_k\}, j \in B$, then

$$\begin{aligned} D_1 &= \sum_{i \in B, j \in C} [(d(i) + d(j))R(i, j) - (d'(i) + d'(j))R'(i, j)] > 0 \\ D_2 &= \left(\sum_{i \in A, j \in B} + \sum_{i \in A, j \in C} \right) [(d(i) + d(j))R(i, j) - (d'(i) + d'(j))R'(i, j)] \\ &= \sum_{j \in B} k(d(u_k) - 1) + \sum_{i \in A \setminus \{u_0, u_k\}, j \in C} (2 + d(j))(k - 2i) + \sum_{j \in C} [k(d(u_0) + d(j)) - (d(u_k) - 1)R(uk, j)] \\ &\quad + \sum_{j \in C} [(d(u_k) + d(j))R(u_k, j) - (1 + d(j))(k + R(u_k, j))] \\ &= |B|k(d(u_k) - 1) + \left(\sum_{i \in A \setminus \{u_0, u_k\}} (k - 2i) \right) \left(\sum_{j \in C} (2 + d(j)) \right) + \sum_{j \in C} k(d(u_0) - 1) \\ &= |B|k(d(u_k) - 1) + |C|k(d(u_0) - 1) > 0. \end{aligned}$$

Therefore, $D_R(G) - D_R(G') = D_1 + D_2 > 0$, the result follows. □

Lemma 2.7. [10] *Let u be a vertex of G such that there are p pendent vertices u_1, u_2, \dots, u_p attached to u . Let v be another vertex of G such that there are q pendent vertices v_1, v_2, \dots, v attached*

to v . Let $G_1 = G - \{vv_1, vv_2, \dots, vv_q\} + \{uv_1, uv_2, \dots, uv_q\}$ and $G_2 = G - \{uu_1, uu_2, \dots, uu_p\} + \{vu_1, vu_2, \dots, vu_p\}$. Then either $D_R(G) > D_R(G_1)$ or $D_R(G) > D_R(G_2)$.

As usual, let P_n and S_n denote the path and star on n vertices. By direct calculation:

$$D_R(S_n) = (n - 1)(3n - 4), \quad D_R(P_n) = \frac{1}{3}n(n - 1)(2n - 1).$$

Let T_n be a tree of order n . In view of Lemma 2.3, Lemma 2.4 and Lemma 2.6, the following corollary immediately implies.

Corollary 2.8. *Among trees of order n , the path P_n has greatest and the star S_n has smallest degree resistance distance. i.e. $3n^2 - 7n + 4 \leq D_R(T_n) \leq \frac{1}{3}n(n - 1)(2n + 1)$.*

A spider is a tree with at most one vertex of degree more than 2, and this vertex is called the hub of the spider (if no vertex of degree more than two, then any vertex can be the hub). A leg of a spider is a path from the hub to a leaf.

Lemma 2.9. *Let $T_{P_n,k}$ be a spider with k legs, and each leg is isomorphic to P_n , then*

$$D_R(T_{P_n,k}) = \frac{1}{3}k(n - 1)(2n - 1)[3(n - 1)(k - 1) + n].$$

Proof. First of all, it is claimed that $D_R(T_{P_n,k}) = kD_R(P_n) + 2mk(k-1)Kf_v(P_n) + k(k - 1)(n - 1)D_v(P_n)$, where $m = n - 1$ is the number of edges in P_n .

By induction on k . If $k = 2$, by Lemma 2.2, calculation shows that $D_R(T_{P_n,2}) = 2D_R(P_n) + 4mKf_v(P_n) + 2(n - 1)D_v(P_n)$, the result holds.

We assume that the equality holds for $k - 1$. By Lemma 2.2 and the induction hypothesis, we get

$$\begin{aligned} D_R(T_{P_n,k}) &= D_R(T_{P_n,k-1}) + D_R(P_n) + 2m(k - 1)Kf_v(P_n) + 2mKf_v(T_{P_n,k-1}) \\ &\quad + (k - 1)(n - 1)D_v(P_n) + (n - 1)D_v(T_{P_n,k-1}) \\ &= (k - 1)D_R(P_n) + 2m(k - 1)(k - 2)Kf_v(P_n) + (k - 1)(k - 2)(n - 1)D_v(P_n) + D_R(P_n) \\ &\quad + 4m(k - 1)Kf_v(P_n) + 2(n - 1)(k - 1)D_v(P_n) \\ &= kD_R(P_n) + 2mk(k - 1)Kf_v(P_n) + k(k - 1)(n - 1)D_v(P_n) \end{aligned}$$

Since $Kf_v(P_n) = \frac{n(n-1)}{2}$ and $D_v(P_n) = (n - 1)^2$, applying Corollary 2.8, we have

$$D_R(T_{P_n,k}) = \frac{1}{3}k(n - 1)(2n - 1)[3(n - 1)(k - 1) + n].$$

□

3. Degree resistance distance of trees with given parameters

Let $\mathcal{T}_{n,d}$ denote the set of all n -vertex trees of diameter d , and $D_{n,d}$ be the n -vertex tree obtained by attaching the center of the star S_{n-d} to the vertex v_l ($l = \lfloor \frac{d}{2} \rfloor + 1$ or $\lceil \frac{d}{2} \rceil + 1$) of a path on $d + 1$ vertices, as shown in Fig. 5.

Theorem 3.1. Among the graphs in $\mathcal{T}_{n,d}$, the tree $D_{n,d}$ is the unique graph with the minimum degree resistance distance.




FIGURE 5. The graphs $D_{n,d}$ and $A_{n,m}$

Proof. Let T be a tree in $\mathcal{T}_{n,d}$ with minimum degree resistance distance. Choose a longest path P in T , say $P = v_1v_2 \cdots v_dv_{d+1}$. It can be confirmed that T is formed by attaching $n - d - 1$ pendant edges to the vertices v_2, v_3, \dots, v_d . If not so, there exists at least one pendant edge uw such that the vertices u, w are distinguished from v_2, v_3, \dots, v_d . Let v be the neighbor of u different from w in T , $T' = T - uw + vw$. Note that $T' \in \mathcal{T}_{n,d}$, applying Lemma 2.3, we can get $D_R(T') < D_R(T)$. It is a contradiction.

Furthermore, we can verify that the $n - d - 1$ pendant edges are all attached to the vertex v_l , that is, the tree T is $D_{n,d}$, where $l = \lfloor \frac{d}{2} \rfloor + 1$ or $l = \lceil \frac{d}{2} \rceil + 1$. If not so, we can choose a vertex $v_j (j \neq \lfloor \frac{d}{2} \rfloor + 1, \lceil \frac{d}{2} \rceil + 1)$ in $V(P)$ such that $d(v_j) \geq 3$ and the path connecting v_j and v_1 or v_{d+1} is a pendant path. Without loss of generality, let the path between v_j and v_1 be a pendant path and $d(v_j, v_1) < d(v_{j+1}, v_{d+1})$. We construct a tree T'' by deleting the pendant edges of v_j (except v_jv_1 if $j = 2$) and adding those edges to the vertex v_{j+1} , obviously, $T'' \in \mathcal{T}_{n,d}$. By lemma 2.5, it implies that $D_R(T'') < D_R(T)$. This is a contradiction. \square

Applying Lemma 2.2 and Corollary 2.8, by calculation we obtain $D_R(D_{n,d})$ as follows:

- (a) $D_R(D_{n,d}) = \frac{1}{3}[9n^2 + (3d^2 - 6d - 21)n - (d^3 + 3d^2 - 10d) + 12]$, if d is even
- (b) $D_R(D_{n,d}) = \frac{1}{3}[9n^2 + (3d^2 - 6d - 18)n - (d^3 + 3d^2 - 7d) + 9]$, if d is odd.

Furthermore, according to Lemma 2.4 and Theorem 3.1, we have the following orders:

$$(3.1) \quad D_R(S_n) = D_R(D_{n,2}) < D_R(D_{n,3}) < \cdots < D_R(D_{n,n-1}) = D_R(P_n).$$

Assume that $\mathcal{P}(n, k)$ is the set of all n -vertex trees with k ($2 \leq k \leq n - 1$) leaves. Let $H_{n,k}$ be a spider with n vertices and k legs, and the lengths of $n - 1 - kr$ legs among them are $r + 1$, the others are r , where $r = \lfloor \frac{n-1}{k} \rfloor$. It is clear that $H_{n,k} \in \mathcal{P}(n, k)$, we get the following result.

Theorem 3.2. Among the graphs in $\mathcal{P}(n, k)$, the tree $H_{n,k}$ has smallest degree resistance distance.

Proof. Let T be a tree in $\mathcal{P}(n, k)$ with minimum degree resistance distance. If $k = 2$ or $n - 1$, the result follows immediately. Therefore, we suppose that $3 \leq k \leq n - 2$.

First of all, we can confirm that T is a spider, that is, T is a tree with at most one vertex of degree more than 2. Otherwise, suppose that there exists two vertices u, v in T such that $d(u) \geq 3$ and $d(v) \geq 3$. Choose a vertex from u, v such that the length of one of its pendent paths is as small as possible. Without loss of generality, let u be the vertex, and $P_u = ux_1 \cdots x$ be the pendent path. Denote by y the neighbor of u in T lying on the path connecting u and v ($y = v$ if u and v are adjacent). Obviously, $d_T(u, x) < \max\{d_T(y, w) : w \in T_v\}$, where T_v is the component of $T - y$ including the vertex v . According to Lemma 2.5, we can obtain a tree $T' \in \mathcal{P}(n, k)$ such that $D_R(T') < D_R(T)$, a contradiction.

Secondly, we can claim that $T \cong H_{n,k}$. Otherwise, in view of Lemma 2.4, it follows that $D_R(T) < D_R(H_{n,k})$, it is a contradiction.

This completes the proof. □

Proposition 3.3. $D_R(H_{n,k}) = \frac{kr}{3}(2r + 1)(3rk - 2r + 1) + \frac{n-kr-1}{2}(5kr^2 - 6r^2 + 3kr + 4rn - 8r + 4n - 4)$, where $r = \lfloor \frac{n-1}{k} \rfloor$.

Proof. Let A be the set of pendent vertices with length of legs $r + 1$ in $H_{n,k}$, $H' = H_{n,k} - A$. By Lemma 2.9, then

$$\begin{aligned} D_R(H_{n,k}) &= \left(\sum_{i,j \in H'} + \sum_{i,j \in A} + \sum_{i \in A, j \in H'} \right) (d(i) + d(j))R(i, j) \\ &= D_R(T_{P_{r+1,k}}) + 2(r + 1)(n - kr - 1)(n - kr - 2) + \\ &\quad (n - kr - 1) \sum_{j \in H'} (1 + d(j))R(i_0, j) (i_0 \in A) \\ &= \frac{kr}{3}(2r + 1)(3rk - 2r + 1) + \frac{n - kr - 1}{2}(5kr^2 - 6r^2 + 3kr + 4rn - 8r + 4n - 4) \end{aligned}$$

□

According to Lemma 2.2 and Lemma 2.4, we get the following orders:

$$(3.2) \quad D_R(S_n) = D_R(H_{n,n-1}) < D_R(H_{n,n-2}) < \cdots < D_R(H_{n,2}) = D_R(P_n).$$

Given a graph G , a subset S of $V(G)$ is called an independent set of G if no pair of vertices of S is adjacent. The independence number $\alpha(G)$ of G is the number of vertices in the largest independent set of G . Recall that if T is a tree of order n , then $\lfloor \frac{n}{2} \rfloor \leq \alpha(T) \leq n - 1$ [4].

Let $\mathcal{Q}_{n,\alpha}$ be the set of n -vertex trees with independent number α . Denote by $I_{n,\alpha}$ the spider obtained from the star $S_{\alpha+1}$ by attaching to $n - \alpha - 1$ of its pendent vertices a new pendent vertex. Obviously, $I_{n,\alpha} \in \mathcal{Q}_{n,\alpha}$.

Theorem 3.4. *Among the graphs in $\mathcal{Q}_{n,\alpha}$, the tree $I_{n,\alpha}$ has smallest degree resistance distance.*

Proof. Let $T \in \mathcal{Q}_{n,\alpha}$, and S be a largest independent set of T with $|S| = \alpha$. Let q be the number of pendent vertices in T , therefore $T \in \mathcal{P}(n, q)$. By Proposition 3.3, we have

$$(3.3) \quad D_R(T) \geq D_R(H_{n,q})$$

For the tree T , one can easily see that $q \leq \alpha$. If $q = \alpha$, considering the fact that $\lfloor \frac{n}{2} \rfloor \leq \alpha \leq n - 1$, then $H_{n,\alpha} \cong I_{n,\alpha}$. If $q < \alpha$, applying Eq. (3.2), then we get

$$(3.4) \quad D_R(H_{n,q}) > D_R(H_{n,q+1}) > \cdots > D_R(H_{n,\alpha}) = D_R(I_{n,\alpha})$$

According to (3.3) and (3.4), it follows that $D_R(T) \geq D_R(I_{n,\alpha})$. This completes the proof. \square

Corollary 3.5. *If $T \in \mathcal{Q}_{n,\alpha}$, then $D_R(T) \geq 4n^2 - \alpha^2 + 4\alpha - 13n + 9$ with equality if and only if $T \cong I_{n,\alpha}$.*

Proof. Since $\lfloor \frac{n}{2} \rfloor \leq \alpha \leq n - 1$, then $r = \lfloor \frac{n-1}{\alpha} \rfloor = 1$. According to Theorem 3.4, we have

$$D_R(I_{n,\alpha}) = D_R(H_{n,\alpha}) = 4n^2 - \alpha^2 + 4\alpha - 13n + 9.$$

By Corollary 3.5, the result follows. \square

By using Lemma 2.3, we have the following conclusion.

Corollary 3.6. *For $\lfloor \frac{n}{2} \rfloor \leq \alpha \leq n - 1$, $D_R(I_{n,\alpha}) < D_R(I_{n,\alpha+1})$.*

A covering of a graph G is a vertex subset $K \subseteq V(G)$ such that each edge of G has at least one end in the set K . The number of vertices in a minimum covering of a graph G is called the covering number of G . Let $\mathcal{K}_{n,m}$ be the set of all n -vertex trees with covering number m , $A_{n,m}$ the tree obtained from the star S_{n-m+1} by attaching $m - 1$ pendent edges to the pendent vertices v_1, v_2, \dots, v_{m-1} of S_{n-m+1} , as shown in Fig.5. It is easy to see that $A_{n,m} \in \mathcal{K}_{n,m}$, where $1 \leq m \leq \lfloor \frac{n}{2} \rfloor$.

Theorem 3.7. [4] *For a bipartite graph G , the sum of the independence number and the covering number of G is equal to the number of vertices of G .*

Theorem 3.8. *Among the graphs in $\mathcal{K}_{n,m}$, the tree $A_{n,m}$ has smallest degree resistance distance.*

Proof. Let $T' \in \mathcal{K}_{n,m}$, and K be a minimum covering set of T' with $|K| = m$. From Theorem 3.8, it follows that the the independence number of T' is equal to $n - m$, then $T' \in \mathcal{Q}_{n,n-m}$. By Corollary 3.5, we have

$$D_R(T') \geq D_R(I_{n,n-m}).$$

By the definition of $A_{n,m}$, $I_{n,n-m}$, we get $A_{n,m} \cong I_{n,n-m}$, thus $D_R(T') \geq D_R(A_{n,m})$. The result follows. \square

In view of Corollary 3.6 and Corollary 3.9, we have the following result.

Corollary 3.9. *If $T \in \mathcal{K}_{n,m}$, then $D_R(T) \geq 3n^2 - m^2 + 2nm - 4m - 9n + 9$ with equality if and only if $T \cong A_{n,m}$.*

A subset M of $E(G)$ is called a matching of G if no two edges in M are adjacent in G . The matching number of a graph G is the maximum cardinality of a matching of G . Denote by $\mathcal{M}_{n,p}$ the set of all n -vertex trees with matching number p .

Theorem 3.10. [4] *In any bipartite graph, the number of edges in a maximum matching is equal to the number of vertices in a minimum covering.*

According to Corollary 3.9 and Corollary 3.11, we obtain the following corollary.

Corollary 3.11. *Among the graphs in $\mathcal{M}_{n,p}$, the tree $A_{n,p}$ is the unique graph with minimum degree resistance distance.*

Let the path $P_n = v_1 v_2 \cdots v_n$, $T(a, n, b)$ be the tree obtained from P_n by attaching a and b pendent vertices to the end vertices v_1 and v_n , respectively. It is clear that the graph $T_{(a,2,b)}$ is the double star $S(a, b)$. We denote by $\mathcal{N}_{n,\Delta}$ the set of all n -vertex trees with maximum degree Δ .

Theorem 3.12. *If $T \in \mathcal{N}_{n,\Delta}$, and $\Delta \geq \frac{n}{2}$, then $D_R(T) \geq 3n^2 - 4\Delta^2 + 4n\Delta - 11n + 8$ with equality if and only if $T \cong T_{(\Delta-1,2,n-\Delta-1)}$.*

Proof. Let T' be the graph with minimum degree resistance distance in $\mathcal{N}_{n,\Delta}$. Let $v \in T'$ and $d(v) = \Delta$. Considering that $\Delta \geq \frac{n}{2}$, T' must satisfy the following three conditions:

- (i) By Lemma 2.3 and Lemma 2.4, T' contains no pendent path with length greater than 1.
- (ii) By Lemma 2.7, All pendent edges of T' are incident to v and another common vertex.
- (iii) By Lemma 2.6, T' contains no internal path with length greater than 1.

Therefore $T' \cong T_{(\Delta-1,2,n-\Delta-1)}$. By calculation, we get $D_R(T') = 3n^2 - 4\Delta^2 + 4n\Delta - 11n + 8$. □

Theorem 3.13. *If $T \in \mathcal{N}_{n,\Delta}$, then $D_R(T) \leq \frac{1}{3}(n-\Delta)(2n\Delta+2n^2-4\Delta^2+15\Delta-3n-11)+(\Delta-1)(3\Delta-4)$ with equality if and only if $T \cong T(\Delta - 1, n - \Delta, 1)$.*

Proof. Choose T' in $\mathcal{N}_{n,\Delta}$, such that its degree resistance distance is as large as possible, and let $v_0 \in T'$ with $d_{T'}(v_0) = \Delta$. In view of Lemma 2.3 and 2.6, any path with end vertex v_0 in T' is a pendent path, that is, $d_{T'}(x) \leq 2$ for any $x \in V(T') \setminus v_0$. According to Lemma 2.4, there are $\Delta - 1$ pendent vertices attached to v_0 , thus $T' \cong T_{(\Delta-1,n-\Delta,1)}$. By direct calculation $D_R(T')$, we get the value, and the result follows. □

Theorem 3.14. *Let T be a n -vertex tree with two vertices of maximum degree Δ , then $D_R(T) \geq D_R(T_{(\Delta-1,n-2\Delta+2,\Delta-1)})$ with equality if and only if $T \cong T_{(\Delta-1,n-2\Delta+2,\Delta-1)}$.*

Proof. Assume that $\mathcal{N}_{n,2\Delta}$ is the set of all n -vertex trees with two vertices of maximum degree Δ . Let $T' \in \mathcal{N}_{n,2\Delta}$ be the tree with maximum degree resistance distance, and let $d_{T'}(v) = d_{T'}(u) = \Delta$. It can be confirmed that $d_{T'}(x) \leq 2$ for any $x \in V(T') \setminus \{v, u\}$, that is, all paths are pendent paths except that between u and v . Otherwise, Let $H \subseteq V(T') \setminus \{v, u\}$ be the set of vertices of T' with degree at least three. There exists a vertex y in H such that y is the end vertex of a pendent path, according to Lemma 2.6, we can get a tree T'' with $D_R(T'') > D_R(T')$, a contradiction.

On the other hand, we show that all pendent paths in T' are pendent edges. Otherwise, choose a shortest pendent path (except pendent edge), without loss of generality, say $vv_1v_2 \cdots v_t$. Move the other pendent paths with the end vertex v from v to v_1 , by Lemma 2.5, we get a tree in $\mathcal{N}_{n,2\Delta}$ such

that the degree resistance distance of the tree is greater than that of T' , it is contradictory to the choice of T' . Thus, $T' \cong T_{(\Delta-1, n-2\Delta+2, \Delta-1)}$. \square

Remark 1: With regard to the problem to determine the trees with maximum degree resistance distance for the given parameters such as number of pendent vertices or independence number, our approach may need to be modified, it would be interesting to continue studying the extremal graphs.

Remark 2: According to (1.1), for the given parameters mentioned above, the extremal graphs corresponding to degree distance, Wiener index and Kirchhoff index among trees are the same as that of DR-index.

REFERENCES

- [1] P. Ali, S. Mukwembi and S. Munyira, Degree distance and vertex-connectivity, *Discrete Appl. Math.*, **161** (2013) 2802–2811.
- [2] C. Arauz, The Kirchhoff indexes of some composite networks, *Discrete Appl. Math.*, **160** (2012) 1429–1440.
- [3] B. Bollobas, *Modern Graph Theory*, Springer-Verlag, 1998.
- [4] J. A. Bondy and U. S. R. Murty, *Graph Theory with Applications*, American Elsevier Publishing Co., Inc., New York, 1976.
- [5] S. B. Chen, Q. Chen, X. Cai and Z. Guo, Maximal Degree Resistance Distance of Unicyclic Graphs, *MATCH Commun. Math. Comput. Chem.*, **75** (2016) 157–168.
- [6] A. A. Dobrynin, R. Entringer and I. Gutman, Wiener index of trees: theory and applications, *Acta Appl. Math.*, **66** (2001) 211–249.
- [7] A. A. Dobrynin, I. Gutman, S. Klavzar and P. Zigert, Wiener index of hexagonal systems, *Acta Appl. Math.*, **72** (2002) 247–294.
- [8] A. A. Dobrynin and A. A. Kochetova, Degree distance of a graph: a degree analogue of the Wiener index, *J. Chem. Inf. Comput. Sci.*, **34** (1994) 1082–1086.
- [9] P. Dankelmann, I. Gutman, S. Mukwembi and H. C. Swart, On the degree distance of a graph, *Discrete Appl. Math.*, **157** (2009) 2773–2777.
- [10] J. F. Du, G. F. Su, J. H. Tu and I. Gutman, The degree resistance distance of cacti, *Discrete Appl. Math.*, **188** (2015) 16–24.
- [11] Z. B. Du and B. Zhou, The Estrada index of trees, *Linear Algebra Appl.*, **435** (2011) 2462–2467.
- [12] L. Feng, G. Yu, K. Xu and Z. Jiang, A note on the Kirchhoff index of bicyclic graphs, *Ars Comb.*, **114** (2014) 33–40.
- [13] I. Gutman, J. Rada and O. Araujo, The Wiener index of starlike trees and a related partial order, *MATCH Commun. Math. Comput. Chem.*, **42** (2000) 145–154.
- [14] I. Gutman, L. Feng and G. Yu, On the degree resistance distance of unicyclic graphs, *Trans. Comb.*, **1** no. 2 (2012) 27–40.
- [15] Q. Guo, H. Deng and D. Chen, The extremal Kirchhoff index of a class of unicyclic graphs, *MATCH Commun. Math. Comput. Chem.*, **61** (2009) 713–722.
- [16] F. Huang, X. l. Li and S. j. Wang, On Maximum Laplacian Estrada Indices of Trees with Some Given Parameters, *MATCH Commun. Math. Comput. Chem.*, **74** (2015) 419–429.
- [17] F. Huang, X. l. Li and S. j. Wang, On maximum Estrada indices of bipartite graphs with some given parameters, *Linear Algebra Appl.*, **465** (2015) 283–295.

- [18] A. Ilic, S. Klavzar and D. Stevanovic, Calculating the degree distance of partial hamming graphs, *MATCH Commun. Math. Comput. Chem.*, **63** (2010) 411–424.
- [19] A. Ilic, A Note on the Additive Degree Kirchhoff Index, *MATCH Commun. Math. Comput. Chem.*, **75** (2016) 223–226.
- [20] D. J. Klein, Resistance-distance sum rules, *Croat. Chem. Acta.*, **75** (2002) 633–649.
- [21] D. J. Klein and M. Randic, Resistance distance, *J. Math. Chem.*, **12** (1993) 81–95.
- [22] S. C. Li, Y. B. Song and H. H. Zhang, On the degree distance of unicyclic graphs with given matching number, *Graphs Combin.*, **31** (2015) 2261–2274.
- [23] J. B. Liu et al. On degree resistance distance of cacti, *Discrete Appl. Math.*, **203** (2016) 217–225.
- [24] M. H. Liu and B. L. Liu, A Survey on Recent Results of Variable Wiener Index, *MATCH Commun. Math. Comput. Chem.*, **69** (2013) 491–520.
- [25] J. L. Palacios, Upper and lower bounds for the additive degree Kirchhoff index, *MATCH Commun. Math. Comput. Chem.*, **70** (2013) 651–655.
- [26] I. Tomescu, Properties of connected graphs having minimum degree distance, *Discrete Math.*, **309** (2009) 2745–2748.
- [27] I. Tomescu, Ordering connected graphs having small degree distances, *Discrete Appl. Math.*, **158** (2010) 1714–1717.
- [28] J. H. Tu, J. f. Du and G. f. Su, The unicyclic graphs with maximum degree resistance distance, *Appl. Math. Comput.*, **268** (2015) 859–864.
- [29] H. Wiener, Structural determination of paraffin boiling points, *J. Am. Chem. Soc.*, **69** (1947) 17–20.
- [30] W. H. Wang, Estrada Indices of the Trees with a Perfect Matching, *MATCH Commun. Math. Comput. Chem.*, **75** (2016) 373–383.
- [31] Y. Yang, X. Jiang, Unicyclic graphs with extremal Kirchhoff index, *MATCH Commun. Math. Comput. Chem.*, **60** (2008) 107–120.
- [32] Y. Yang, D.J. Klein, Resistance distance-based graph invariants of subdivisions and triangulations of graphs, *Discrete Appl. Math.*, **181** (2015) 260–274.
- [33] B. Zhou, N. Trinajstić, On resistance-distance and kirchhoff index, *J. Math. Chem.*, **46** (2009) 283–289.
- [34] H. Zhang, X. Jiang, Y. Yang, Bicyclic graphs with extremal Kirchhoff index, *MATCH Commun. Math. Comput. Chem.*, **61** (2009) 697–712.

Fanguo He

College of Mathematics and Physics, Huanggang Normal University, Huanggang, China

Email: hfg0118@126.com

Xinnong Jiang

College of Life Science and Technology, Huazhong University of Science and Technology, Wuhan, China

Email: jiangxn88@hust.edu.cn