



www.combinatorics.ir

---

**Transactions on Combinatorics**

ISSN (print): 2251-8657, ISSN (on-line): 2251-8665

Vol. 4 No. 2 (2015), pp. 23-35.

© 2015 University of Isfahan

---



www.ui.ac.ir

## THE SECOND IMMANANT OF SOME COMBINATORIAL MATRICES

R. B. BAPAT AND S. SIVASUBRAMANIAN\*

Communicated by Alireza Abdollahi

**ABSTRACT.** Let  $A = (a_{i,j})_{1 \leq i,j \leq n}$  be an  $n \times n$  matrix where  $n \geq 2$ . Let  $\det_2(A)$ , its second immanant be the immanant corresponding to the partition  $\lambda_2 = 2, 1^{n-2}$ . Let  $G$  be a connected graph with blocks  $B_1, B_2, \dots, B_p$  and with  $q$ -exponential distance matrix  $\text{ED}_G$ . We give an explicit formula for  $\det_2(\text{ED}_G)$  which shows that  $\det_2(\text{ED}_G)$  is independent of the manner in which  $G$ 's blocks are connected. Our result is similar in form to the result of Graham, Hoffman and Hosoya and in spirit to that of Bapat, Lal and Pati who show that  $\det \text{ED}_T$  where  $T$  is a tree is independent of the structure of  $T$  and only dependent on its number of vertices. Our result extends more generally to a product distance matrix associated to a connected graph  $G$ . Similar results are shown for the  $q$ -analogue of  $T$ 's laplacian and a suitably defined matrix for arbitrary connected graphs.

### 1. Introduction

We consider the second immanant of  $n \times n$  matrices with entries from a commutative ring. We briefly state some needed background from the representation theory of the symmetric group  $\mathfrak{S}_n$  on the set  $[n] = \{1, 2, \dots, n\}$  over  $\mathbb{C}$ , the complex numbers. Let  $A = (a_{i,j})_{1 \leq i,j \leq n}$  be an  $n \times n$  matrix with entries from a commutative ring. In this work, all matrices will be either over  $\mathbb{Z}$ , the integers or over the polynomial ring  $\mathbb{R}[q]$  where  $q$  is a variable. Let  $f : \mathfrak{S}_n \rightarrow \mathbb{Z}$  be a function. Define  $\det_f(A) = \sum_{\sigma \in \mathfrak{S}_n} f(\sigma) \prod_{i=1}^n a_{i,\sigma(i)}$ . We only consider functions  $f : \mathfrak{S}_n \mapsto \mathbb{Z}$  that arise as characters of irreducible representations of  $\mathfrak{S}_n$  over  $\mathbb{C}$ . If  $f$  is such a function, we call  $\det_f(A)$  as an immanant. When  $f$  is the  $\text{sgn}$  function defined as  $f(\pi) = \text{sgn}(\pi)$  for all  $\pi \in \mathfrak{S}_n$ , then, clearly  $\det_{\text{sgn}}(A)$  is the usual determinant of  $A$ . That is, we have  $\det_{\text{sgn}}(A) = \det A$ . If  $f$  is the id or all ones function defined as  $f(\pi) = 1$  for all  $\pi \in \mathfrak{S}_n$ , then  $\det_{\text{id}}(A) = \text{perm}(A)$ , where  $\text{perm}(A)$  is the permanent of  $A$ .

---

MSC(2010): Primary: 15A15; Secondary: 05C05.

Keywords: Immanant, Product-distance matrix, Laplacian.

Received: 20 February 2014, Accepted: 10 September 2014.

\*Corresponding author.

The number of distinct irreducible representations of  $\mathfrak{S}_n$  is  $p(n)$ , the number of partitions of the positive integer  $n$  (see Sagan’s book [10, Proposition 1.10.1]). Thus, if  $\lambda$  is a partition of  $n$ , we have functions  $\chi_\lambda : \mathfrak{S}_n \rightarrow \mathbb{Z}$ . As seen, both the determinant and the permanent of a matrix are immanants, with the determinant corresponding to the partition  $(1, 1, \dots, 1)$  and the permanent corresponding to the partition  $(n)$ .

Let  $n \geq 2$  and  $\lambda_2$  be the partition  $(2, 1^{n-2})$  of  $n$ . Denote as  $\chi_2 : \mathfrak{S}_n \rightarrow \mathbb{Z}$ , the irreducible character of  $\mathfrak{S}_n$  corresponding to the partition  $\lambda_2$ . Define the second immanant of  $A$  to be

$$\det 2(A) = \sum_{\pi \in \mathfrak{S}_n} \chi_2(\pi) \prod_{i=1}^n a_{i, \pi(i)}.$$

For an  $n \times n$  matrix  $A$ ,  $\det 2(A)$  can be computed efficiently. Littlewood’s book [7, Chapter 6.5] contains a nice exposition of this result. See the work of Merris and Watkins [9] as well. For an  $n \times n$  matrix  $A$  and for  $1 \leq i \leq n$ , let  $A(i)$  be the  $(n - 1) \times (n - 1)$  matrix obtained from  $A$  by deleting its  $i$ -th row and its  $i$ -th column.

**Theorem 1.1.** *Let  $A = (a_{i,j})_{1 \leq i,j \leq n}$  be an  $n \times n$  matrix. Then,*

$$\det 2(A) = \sum_{i=1}^n a_{i,i} \det A(i) - \det A.$$

We now move on to the matrices we consider in this work. Let  $G$  be an undirected, connected graph on the vertex set  $V(G) = [n]$ . A *block* of  $G$  is a maximally connected subgraph without a cut-vertex. For a graph  $G$ , we will look at functions  $\eta : V(G) \times V(G) \rightarrow R$  where  $R$  is a commutative ring (either  $\mathbb{Z}$  or  $\mathbb{R}[q]$ ). A *product distance* on  $G$  is a function  $\eta : V(G) \times V(G) \rightarrow R$ , that satisfies the following two conditions:

- (1)  $\eta(i, i) = 1$  for all  $i \in [n]$ .
- (2) if  $i, j \in V(G)$  are vertices such that every path from  $i$  to  $j$  passes through the cut-vertex  $k$ , then  $\eta(i, j) = \eta(i, k)\eta(k, j)$ .

We emphasize that we start with an undirected, connected graph and for each of its block, assign distances to pairs of vertices within the block. Thus, we essentially have the freedom to assign *distances* within each block subject to diagonal entries being 1. Rule 2 (the product rule) is then used to obtain distances between pairs of vertices from different blocks. Because of this *product rule*, we call such distances as *product distances* and use the terms distance and product distance interchangeably. Since we need not have symmetric distances within each block, it is not necessary that  $\eta(i, j) = \eta(j, i)$  for  $i, j \in V(G)$ . As distances could be asymmetric, we could alternatively phrase our results using the terminology of directed graphs. We prefer to use undirected graph terminology in this work.

We will form a matrix with entries being  $\eta(i, j)$  and sometimes write  $\eta_{i,j}$  alternatively to denote  $\eta(i, j)$ . Let  $G$  have blocks  $H_1, H_2, \dots, H_r$ . Let  $\eta(\cdot, \cdot)$  be a product distance on  $G$  and let  $D_G = (\eta_{i,j})_{1 \leq i,j \leq n}$  be the corresponding distance matrix.

The definition of product-distances is motivated by a concrete example: the exponential distance matrix of a connected graph  $ED_G$ . Given a connected graph  $G$  on the vertex set  $[n]$ , let the distance

between two vertices  $i, j \in V(G)$  be denoted  $d_{i,j}$ . i.e.  $d_{i,j}$  is the length of the minimum length path from  $i$  to  $j$  in  $G$ . Define the  $n \times n$  matrix  $ED_G = (q^{d_{i,j}})_{1 \leq i, j \leq n}$  as the exponential distance matrix where  $q$  is an indeterminate and  $q^0 = 1$ . It can be readily checked that  $\eta(i, j) = q^{d_{i,j}}$  is a product distance that is symmetric. If  $\eta(\cdot, \cdot)$  is a product distance on  $G$  and if  $G$  has blocks  $H_1, H_2, \dots, H_r$ , then, each  $H_i$  is a graph in its own right and thus has an induced product distance matrix  $D_{H_i}$  obtained by restricting  $\eta(\cdot, \cdot)$  to pairs of vertices, both in  $H_i$ . If the graph  $G$  is clear from the context, we abridge  $D_G$  to  $D$ .

If  $D$  is a matrix whose entries form a product distance on  $G$ , Bapat and Sivasubramanian [2] showed that  $\det D$  only depends on  $\det D_{H_i}$  for individual blocks  $H_i$  of  $G$  and not on the manner in which the  $H_i$ 's are connected.

**Theorem 1.2.** ([2, Theorem 4] ) *Let  $G$  be a connected graph with blocks  $H_i, 1 \leq i \leq r$  and product distance matrix  $D_G$ . For each such  $i$ , let the product distance matrix of each  $H_i$  be  $D_{H_i}$ . Then,*

$$\det D_G = \prod_{i=1}^r \det D_{H_i}$$

In particular,  $\det D_G$  is independent of the manner in which the blocks  $H_i$  of  $G$  are connected. In this work, we extend this result to  $\det 2(D_G)$  by giving an explicit formula for  $\det 2(D_G)$  in terms of the determinant and the second immanant of the  $D_{H_i}$ 's (see Theorem 3.5). Our formula is identical in form to the formula for the determinant of the distance matrix of a connected graph given by Graham, Hoffman and Hosoya. Their formula with the relevant background appears in Subsection 2.2.

As exponential distance matrices are special cases of product distances, considering the case when  $G$  is a tree  $T$ , we get the result that  $\det 2(ED_T)$  is independent of the structure of the tree  $T$ . For this special case when  $G$  is a tree, our result is true in a more general non-commutative setting with *matrix weights* on the edges  $e_i$ . Let  $T$  be a tree with vertex set  $[n]$ . Let the edge  $e_i$  of  $T$  have a *matrix weight*  $W_i$  (for  $1 \leq i < n$ ) and where each  $W_i$  is an  $s \times s$  matrix over a commutative ring  $R$ . For  $i, j \in [n]$ , clearly, there is a unique path  $p_{i,j}$  between  $i$  and  $j$  given by the sequence of edges  $e_1, e_2, \dots, e_r$  where  $i \in e_1, j \in e_r$  and there is a common vertex in the edges  $e_i, e_{i+1}$  for  $1 \leq i < r$ . Define  $d_{i,j}$ , the distance between vertices  $i$  and  $j$  as  $\prod_{i=1}^r W_i$  where the product takes the order of the matrices as they appear in  $p_{i,j}$  into account. Note that  $d_{i,j}$  is an  $s \times s$  matrix. When  $i = j$ , define  $d_{i,j}$  to be the  $s \times s$  identity matrix. Consider the  $ns \times ns$  matrix  $\mathcal{D}_T = (d_{i,j})_{1 \leq i, j \leq n}$  (i.e. we have a block matrix). We call  $\mathcal{D}_T$  as the *non-commutative analogue* of the distance matrix of  $T$ . When the tree  $T$  is clear, we abuse notation and write  $\mathcal{D}$  instead of the more precise  $\mathcal{D}_T$ . Bapat and Sivasubramanian [3, Theorem 3] showed the following.

**Theorem 1.3.** *Let  $T$  be a tree on  $n$  vertices and for  $1 \leq i < n$ , let edge  $e_i$  have an  $s \times s$  matrix weight  $W_i$ . Then,  $\det \mathcal{D}_T = \prod_{i=1}^{n-1} (I - W_i^2)$ . In particular,  $\det \mathcal{D}_T$  is independent of the structure of the tree  $T$  and only depends on  $n$  and the weights  $W_1, W_2, \dots, W_{n-1}$ .*

We show a second-immanant analogue of Theorem 1.3 by giving an explicit formula for  $\det 2(\mathcal{D}_T)$  (see Theorem 3.8). Our proofs rely on explicit inverse results found by Bapat and Sivasubramanian and on Theorem 1.1.

## 2. Second immanant of combinatorial matrices

We consider two families of matrices in this section and give relevant background on their second-immanants. We recall that all graphs in this work are connected.

**2.1. Laplacian matrices.** Let  $G$  be a connected graph with adjacency matrix  $A$  and diagonal matrix  $D$  with  $D(i, i) = \deg(i)$ , where  $\deg(i)$  is the degree of vertex  $i$ . The laplacian of  $G$  is the matrix  $L = D - A$ . The following result of Merris [8] is immediate when we combine Theorem 1.1 with the Matrix Tree Theorem (see West's book [15, Page 86]).

**Corollary 2.1.** *Let  $G$  be a connected graph with  $n$  vertices,  $m$  edges and  $\kappa$  spanning trees. Let  $L$  be its laplacian matrix. Then,  $\det 2(L) = 2m\kappa$ .*

Below, we mention a special case of Corollary 2.1, when the graphs are trees. We mention this as we generalise this case when  $G$  is a tree to the  $q$ -analogue of the laplacian of a tree (see Corollary 4.5).

**Corollary 2.2.** *Let  $T$  be a tree on  $n$  vertices and let  $L$  be its laplacian matrix. Then,  $\det 2(L) = 2(n - 1)$ . Hence,  $\det 2(L)$  only depends on  $n$  and is independent of the structure of the tree  $T$ .*

**2.2. Distance matrices.** Let  $G$  be a connected graph with vertex set  $[n]$  and with distance matrix  $D = (d_{i,j})_{1 \leq i, j \leq n}$ . Thus  $d_{i,j}$  is the length of the shortest path between  $i$  and  $j$  in  $G$  and  $d_{i,i} = 0$  for all  $i \in [n]$ . We begin with the following result of Grone and Merris (see [6, Page 590]).

**Lemma 2.3.** *If  $A = (a_{i,j})_{1 \leq i, j \leq n}$  is an  $n \times n$  matrix such that  $a_{i,i} = 0$  for all  $1 \leq i \leq n$ , then  $\det 2(A) = -\det A$ .*

Thus, for distance matrices  $D$  of connected graphs  $G$ , up to sign  $\det D = \det 2(D)$ . The following is a well known result of Graham and Pollak [5].

**Theorem 2.4** (Graham and Pollak). *Let  $T$  be a tree with vertex set  $[n]$  with distance matrix  $D_T$ . Then,  $\det D_T = (-1)^{n-1}(n - 1)2^{n-2}$ . Thus,  $\det D_T$  only depends on  $n$  and is independent of the structure of the tree  $T$ .*

Later, Graham, Hoffman and Hosoya [4] proved a more general and more attractive theorem about the determinant of the distance matrix  $D_G$  of a strongly connected digraph  $G$  as a function of the distance matrix of its *2-connected blocks* (also called blocks). Denote the sum of the cofactors of a matrix  $A$  as  $\text{cofsum}(A)$ . Graham, Hoffman and Hosoya (see [4]) showed the following.

**Theorem 2.5** (Graham, Hoffman and Hosoya). *Let  $G$  be a strongly connected digraph with 2-connected blocks  $G_1, G_2, \dots, G_r$ . Then,*

- (1)  $\text{cofsum}(D_G) = \prod_{i=1}^r \text{cofsum}(D_{G_i})$  and
- (2)  $\det D_G = \sum_{i=1}^r \det D_{G_i} \prod_{j \neq i} \text{cofsum}(D_{G_j})$ .

Graham, Hoffman and Hosoya’s theorem implies that  $\det D_G$  is independent of the manner in which the blocks of  $G$  are connected. It is also easy to recover Theorem 2.4 from Theorem 2.5 (as all blocks of  $T$  are  $K_2$ , the complete graph on 2 vertices). Since  $D_T$  and  $D_G$  are distance matrices, all their diagonal elements are zero. Further,  $q$ -analogues of Theorems 2.4 and 2.5 were given by Bapat, Lal and Pati [1] and by Sivasubramanian [12] respectively. Here, each positive integer  $n$  is replaced by the polynomial  $[n]_q = 1 + q + q^2 + \dots + q^{n-1}$ , and  $[0]_q = 0$ . Denote the  $q$ -analogues of  $D_T$  and  $D_G$  as  $qD_T$  and  $qD_G$ , respectively. Thus, by Lemma 2.3, we have the following simple corollary.

**Corollary 2.6.** *Let  $G$  be a connected graph on the vertex set  $[n]$  with distance matrix  $D_G$  and with  $qD_G$  being the  $q$ -analogue of  $D_G$ . Then, both  $\det 2(D_G)$  and  $\det 2(qD_G)$  are independent of the tree-like manner of connection of its blocks.*

Consider the case now when  $D_G$  is the product distance matrix of a graph  $G$ . Since all the diagonal elements of  $D_G$  are 1, Lemma 2.3 is not applicable. It is these matrices whose second immanant we find in this work.

### 3. The second immanant of $D_G$ and $D_T$

Let  $G$  be a graph with blocks  $B_1, B_2, \dots, B_p$  and with a product distance  $\eta(\cdot, \cdot)$ . Let  $D_G = (\eta_{i,j})$  be the matrix of product distances on  $G$ . Let the restriction of  $D_G$  to vertices of  $B_i$  be denoted  $D_{B_i}$  for all  $1 \leq i \leq p$ . Assume that for all  $1 \leq i \leq p$ , we have  $\det D_{B_i} \neq 0$ . That is, assume all the matrices  $D_{B_i}$  are invertible. For each  $1 \leq i \leq p$ , let  $D_{B_i}^{-1} = N_i$ . If  $|V(B_i)| = n_i$ , then  $N_i$  has dimension  $n_i \times n_i$ . Let  $M_i$  be the matrix  $N_i$  enlarged to have dimension  $n \times n$  with zeroes added for all entries outside  $V(B_i)$ .




FIGURE 1. Decomposing  $G$  into blocks.




FIGURE 2. The block cut-point graph of  $G$  of Figure 1.

We illustrate getting the matrix  $M_i$  from  $N_i$  on an example. For the graph given in Figure 1, clearly, there are four blocks. These are marked as  $B_1, B_2, B_3$  and  $B_4$  respectively. We show how to get  $M_2$  from  $N_2$ . As shown in the figure,  $B_2$  is the block consisting of the edge  $\{1, 4\}$ . If  $B_2$  has  $N_2$  as given

below, then we obtain  $M_2$  by padding zeroes outside all vertices of  $B_2$ . As the two vertices of  $B_2$  are 1 and 4,  $M_2$  will be as given below.

$$N_2 = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \quad M_2 = \begin{pmatrix} A & 0 & 0 & B & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ C & 0 & 0 & D & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad R = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

Let  $G$  have  $\ell$  cut vertices  $w_1, w_2, \dots, w_\ell$  in between its blocks. In the block cutpoint graph of  $G$  (see West’s book [15, Page 156]), let the degree of  $w_i$  be  $d_i$ . Let  $R$  be the diagonal matrix with its  $(w_i, w_i)$ -th entry being  $d_i - 1$  for each  $1 \leq i \leq \ell$ . For non cut-vertices  $a$ , define the  $(a, a)$ -th entry of  $R$  to be 0. For the graph given in Figure 1, we have its block cutpoint graph given in Figure 2. For this graph, we have  $p = 4$  ( $G$  has four blocks) and  $\ell = 2$  ( $G$  has two cut vertices). Let  $w_1 = 4$  and  $w_2 = 1$  be the cut vertices ( $w_i$  is the index of the  $i$ -th cut-vertex). From the block-cutpoint graph, we see that  $d_1 = 3, d_2 = 2$  (number of blocks incident on the  $w_i$ ’s). The matrix  $R$  for the graph of Figure 1 is given above.

Form an  $n \times n$  matrix  $K = \sum_{i=1}^p M_i - R$ . We reiterate below the steps required to obtain  $K$ .

- (1) For each block  $B_i$ , where  $1 \leq i \leq p$ , form  $M_i$  by padding zeroes to the inverse matrix  $N_i = D_{B_i}^{-1}$  at all indices outside  $B_i$ .
- (2) Form the diagonal matrix  $R$  from the block-cutpoint graph of  $G$ .
- (3) Set  $K = \sum_{i=1}^p M_i - R$ .

With the above definitions, we have the following result of Bapat and Sivasubramanian.

**Theorem 3.1.** ([2, Theorem 6]) *Let  $G$  be a connected graph with blocks  $B_1, B_2, \dots, B_p$ . Let  $D_G$  be the product distance matrix of  $G$  and for  $1 \leq i \leq p$ , let  $D_{B_i}$  be the restriction of  $D$  to the vertices in  $B_i$ . If  $\det D_{B_i} \neq 0$  for all  $1 \leq i \leq p$ , then,  $D_G^{-1} = K$ .*

We begin with the following simple lemma. For a matrix  $A$ , denote its trace as  $\text{Trace}(A)$ .

**Lemma 3.2.** *Let  $A = (a_{i,j})_{1 \leq i,j \leq n}$  be an invertible  $n \times n$  matrix with  $a_{i,i} = 1$  for all  $i$ . Then,*

$$\text{Trace}(A^{-1}) = 1 + \frac{\det 2(A)}{\det A}.$$

**Proof 3.3.** *We use Theorem 1.1. Since  $a_{i,i} = 1$  for all  $i$ ,  $\det 2(A) = \det A \cdot \text{Trace}(A^{-1}) - \det A$ , completing the proof.*

**Remark 3.4.** *Recall the matrices  $N_i$  and  $M_i$  given at the beginning of this section. We will apply Lemma 3.2 to the matrices  $M_i$ . Since  $\text{Trace}(M_i) = \text{Trace}(N_i)$  and the matrix  $N_i = D_{B_i}^{-1}$ , we get  $\text{Trace}(M_i) = 1 + \frac{\det 2(D_{B_i})}{\det D_{B_i}}$ .*

We are now ready to prove our first main result.

**Theorem 3.5.** *Let  $D_G$  be the product distance matrix of a connected graph  $G$  with blocks  $B_1, B_2, \dots, B_p$ , with the property that  $\det D_{B_i} \neq 0$  for all  $1 \leq i \leq p$ . Then,*

$$\det 2(D_G) = \sum_{i=1}^p \left[ \det 2(D_{B_i}) \prod_{j \neq i} \det D_{B_j} \right].$$

*In particular,  $\det 2(D_G)$  is independent of the tree-like manner of connection of  $G$ 's blocks.*

**Proof 3.6.** *Since the  $(i, i)$ -th element of  $D_G$  is 1 for all  $i$ , using Theorems 1.1 and 3.1, we have*

$$\det 2(D_G) = \det D_G \cdot \text{Trace}(K) - \det D_G = \det D_G (\text{Trace}(K) - 1).$$

*By definition, we have  $\text{Trace}(K) = \sum_{i=1}^p \text{Trace}(M_i) - \text{Trace}(R)$ . It is easy to see by induction on the number of blocks in  $G$  that  $\text{Trace}(R) = p - 1$ . Thus,*

$$\begin{aligned} \det 2(D_G) &= \det D_G \left( \sum_{i=1}^p \text{Trace}(M_i) - p \right) = \det D_G \left( \sum_{i=1}^p \left\{ \frac{\det 2(D_{B_i})}{\det D_{B_i}} + 1 \right\} - p \right) \\ &= \det D_G \left( \sum_{i=1}^p \frac{\det 2(D_{B_i})}{\det D_{B_i}} \right) = \sum_{i=1}^p \det 2(D_{B_i}) \prod_{j \neq i} \det D_{B_j}. \end{aligned}$$

*We have used Remark 3.4 in the second line above and Theorem 1.2 in the third line. This completes the proof.*

We note that in Theorem 3.1, we do not need the product distance to be symmetric. i.e. we do not require  $\eta_{i,j} = \eta_{j,i}$ . It is simple to see that symmetry of the product distance is not required for Theorem 3.5 either. Our earlier proof crucially uses the fact that each diagonal entry of  $D_G$  is 1 and our explicit inverse result.

We note that Theorem 3.5 and Theorem 1.2 are for product distance matrices. They are counterparts of Theorem 2.5 for ordinary distance matrices with  $\det$  and  $\det 2()$  playing the roles of  $\text{cofsum}()$  and  $\det$  respectively.

**3.1.  $\det 2(\mathcal{D}_T)$  for a tree  $T$ .** A similar proof gives us a non-commutative analogue of Theorem 3.5 for trees as described in Section 1. Recall  $\mathcal{D}_T$  defined for a tree  $T$  on  $n$  vertices with edge  $e_i$  bearing a matrix weight  $W_i$  for  $1 \leq i < n$ . As each  $W_i$  is an  $s \times s$  matrix,  $\mathcal{D}_T$  is an  $ns \times ns$  matrix. Bapat and Sivasubramanian showed that if for each  $1 \leq i < n$ , the  $s \times s$  matrix  $(I - W_i^2)$  is invertible, then the inverse of  $\mathcal{D}_T$  can be written explicitly. To describe it, recall that if  $A = (a_{i,j})_{1 \leq i \leq m, 1 \leq j \leq n}$  is an  $m \times n$  matrix and  $B$  is a  $p \times q$  matrix, then their Kronecker product  $A \otimes B$  is the  $mp \times nq$  matrix given by

$$A \otimes B = \begin{pmatrix} a_{1,1}B & a_{1,2}B & \cdots & a_{1,n}B \\ a_{2,1}B & a_{2,2}B & \cdots & a_{2,n}B \\ \vdots & \vdots & \ddots & \vdots \\ a_{m,1}B & a_{m,2}B & \cdots & a_{m,n}B \end{pmatrix}$$

Let the degree sequence of  $T$  be  $d_1, d_2, \dots, d_n$  and define an  $n \times n$  diagonal matrix  $\text{Deg}$  by  $\text{Deg} = \text{Diag}(d_1, d_2, \dots, d_n)$ . Define the  $ns \times ns$  matrix  $\Delta = \text{Deg} \otimes I_s$  (thus  $\Delta$  has  $n$  non-zero diagonal blocks, each of size  $s \times s$ ). For an  $s \times s$  matrix  $P$ , if the matrix  $I - P^2$  is invertible, then it is clear that

$$(3.1) \quad \begin{pmatrix} I & P \\ P & I \end{pmatrix}^{-1} = \begin{pmatrix} (I - P^2)^{-1} & -P(I - P^2)^{-1} \\ -P(I - P^2)^{-1} & (I - P^2)^{-1} \end{pmatrix}.$$

Recall that edge  $e_k$  has an  $s \times s$  matrix weight  $W_k$  attached to it. For each edge  $e_k = \{i, j\}$  of  $T$ , consider the following  $ns \times ns$  matrix  $M_{e_k}$  which we describe in terms of  $s \times s$  blocks as follows. The  $(i, i)$ -th and  $(j, j)$ -th blocks of  $M_{e_k}$  are  $(I - W_k^2)^{-1}$  and the  $(i, j)$ -th and  $(j, i)$ -th blocks are  $-W_k(I - W_k^2)^{-1}$ . For other indices  $(a, b)$ , define the  $(a, b)$ -block to be the  $s \times s$  zero block. We have  $n - 1$  such matrices  $M_{e_k}$  one for each edge  $e_k$  where  $1 \leq k \leq n - 1$ . With these definitions, the following inverse result was proved by Bapat and Sivasubramanian [3, Theorem 4].

**Theorem 3.7.** *Let  $\mathcal{D}_T$  be the non-commutative analogue of the distance matrix of a tree  $T$  on  $n$  vertices with edge  $e_k$  having matrix weight  $W_k$  for  $1 \leq k \leq n - 1$ . Then,  $\mathcal{D}_T^{-1} = I - \Delta + \sum_{k=1}^{n-1} M_{e_k}$ .*

With this background, we can now show our next main result.

**Theorem 3.8.** *Let  $T$  be a tree on  $n$  vertices with edge  $e_i$  having an  $s \times s$  matrix weight  $W_i$  and let  $\mathcal{D}_T$  be the non-commutative analogue of its distance matrix. Recall  $I_s$  is the  $s \times s$  identity matrix and for  $1 \leq i < n$ , define the  $2s \times 2s$  matrix  $L_i = \begin{pmatrix} I_s & W_i \\ W_i & I_s \end{pmatrix}$ . Then,*

$$\det 2(\mathcal{D}_T) = \left( \sum_{i=1}^{n-1} \det 2(L_i) \prod_{j \neq i} \det L_j \right) - (n - 2)(s - 1) \det \mathcal{D}_T.$$

*In particular,  $\det 2(\mathcal{D}_T)$  is independent of the structure of  $T$  and only depends on  $n$  and the matrices  $W_i$  for  $1 \leq i < n$ .*

**Proof 3.9.** *Using Theorem 1.1 and the fact that all diagonal entries are 1, we get,  $\det 2(\mathcal{D}_T) = \det \mathcal{D}_T \cdot \text{Trace}(\mathcal{D}_T^{-1}) - \det \mathcal{D}_T$ . By Theorem 3.7,  $\text{Trace}(\mathcal{D}_T^{-1}) = \text{Trace}(I - \Delta + \sum_{k=1}^{n-1} M_{e_k})$ , where all matrices have dimension  $ns \times ns$ . Breaking this up into two terms, we get  $\text{Trace}(I - \Delta) = s(n - (2n - 2)) = s(2 - n)$  and the term  $\text{Trace}(\sum_{k=1}^{n-1} M_{e_k})$ . Recall the  $2s \times 2s$  matrix  $L_k = \begin{pmatrix} I_s & W_k \\ W_k & I_s \end{pmatrix}$ , where all four block matrices are of dimension  $s \times s$ . For all  $1 \leq k < n$ , since  $\text{Trace}(M_{e_k}) = \text{Trace}(L_k^{-1})$  and since  $L_k$  has all diagonal entries 1, by Lemma 3.2, we get that  $\text{Trace}(L_k^{-1}) = \left(1 + \frac{\det 2(L_k)}{\det L_k}\right)$ . Thus,*

$$\begin{aligned} \det 2(\mathcal{D}_T) &= \det \mathcal{D}_T \text{Trace}(\mathcal{D}_T^{-1}) - \det \mathcal{D}_T \\ &= \det \mathcal{D}_T \left( s(2 - n) + \sum_{k=1}^{n-1} \left\{ 1 + \frac{\det 2(L_k)}{\det L_k} \right\} - 1 \right) \\ &= \det \mathcal{D}_T \left( \sum_{k=1}^{n-1} \frac{\det 2(L_k)}{\det L_k} \right) + \det(\mathcal{D}_T)(s - 1)(2 - n) \\ &= \left( \sum_{i=1}^{n-1} \det 2(L_i) \prod_{j \neq i} \det L_j \right) - (n - 2)(s - 1) \det \mathcal{D}_T. \end{aligned}$$



In the last line, we have used Theorem 1.3. The proof is complete.

We note that when each  $W_i$  is the  $1 \times 1$  indeterminate  $w_i$ , then  $L_i = \begin{pmatrix} 1 & w_i \\ w_i & 1 \end{pmatrix}$ . That is,  $L_i$  is a matrix with  $L_i = D_{B_i}$ . It is easy to check that  $\det L_i = 1 - w_i^2$ ,  $\det 2(L_i) = 1 + w_i^2$ . In this case, we recover a special case of Theorem 3.5.

**3.2. Monomial immanant corresponding to  $\lambda = 2, 1^{n-2}$ .** In this subsection, we show that our results can be stated in the language of a monomial immanant. To describe monomial immanants, we need a few preliminaries from the theory of symmetric functions. We refer the reader to Stanley [13, Chapter 7] for relevant background. Given any symmetric function  $f$  of degree  $n$  in infinitely many variables  $x_1, x_2, \dots$ , we can get a function  $\psi_f : \mathfrak{S}_n \rightarrow \mathbb{Z}$  as follows.

Recall that each permutation  $\pi \in \mathfrak{S}_n$  can be written in cycle notation. Let  $\pi$  have  $\ell$ -cycles  $C_1, C_2, \dots, C_\ell$  with  $t_i = |C_i|$  for  $1 \leq i \leq \ell$ . Since there is no order among the cycles  $C_i$  of  $\pi$ , we assume that  $t_1 \geq t_2 \geq \dots \geq t_\ell$ . Thus,  $w(\pi) = (t_1, t_2, \dots, t_\ell)$  is a partition of the positive integer  $n$  and hence for each  $\pi \in \mathfrak{S}_n$ , we get a partition  $w(\pi)$  of the integer  $n$ . We write this as  $w(\pi) \vdash n$ .

Symmetric functions of degree  $n$  with rational coefficients form a vector space denoted  $\Lambda_{\mathbb{Q}}^n$  which is equipped with a standard inner product. To define the inner product, consider the basis,  $\{m_\lambda\}$  for  $\lambda \vdash n$ , of monomial immanants and the basis  $\{h_\mu\}$  for  $\mu \vdash n$ , of complete homogenous symmetric functions. That is, both bases are indexed by partitions  $\lambda \vdash n$ . If  $f, g \in \Lambda_{\mathbb{Q}}^n$ , write  $f = \sum_{\lambda \vdash n} A_\lambda m_\lambda$  and  $g = \sum_{\mu \vdash n} B_\mu h_\mu$ . Their inner-product denoted  $\langle f, g \rangle$  is defined as  $\langle f, g \rangle = \sum_{\lambda \vdash n} A_\lambda B_\lambda$ .

Another basis for degree  $n$  symmetric functions are the power sum symmetric functions  $p_\lambda$  for  $\lambda \vdash n$ . We are now in a position to define the map  $\psi_f : \mathfrak{S}_n \rightarrow \mathbb{Z}$ . Define  $\psi_f : \mathfrak{S}_n \rightarrow \mathbb{Z}$  by  $\psi_f(\pi) = \langle f, p_{w(\pi)} \rangle$ .

Consider the partition  $\lambda_2 = (2, 1^{n-2})$  and the symmetric function  $m_{\lambda_2}$ . Define  $\psi_2 = \psi_{m_{\lambda_2}}$ . Thus  $\psi_2 : \mathfrak{S}_n \rightarrow \mathbb{Z}$  is the function obtained from the monomial symmetric function  $m_{\lambda_2}$ . For an  $n \times n$  matrix  $A$ , consider the immanant  $\det_{\psi_2}(A)$  defined with respect to  $\psi_2$ . That is,  $\det_{\psi_2}(A)$  is the immanant defined with respect to the monomial symmetric function  $m_{\lambda_2}$ . Such immanants are referred to as monomial immanants. We need the following result (see Stembridge, proof of Theorem 2.7 [14]).

**Theorem 3.10.** *Let  $A = (a_{i,j})_{1 \leq i,j \leq n}$  be an  $n \times n$  matrix. Then,*

$$\det_{\psi_2}(A) = \sum_{i=1}^n a_{i,i} \det A(i) - n \det A.$$

We get the following simple corollary from Theorems 3.10 and 1.1.

**Corollary 3.11.** *Let  $A$  be an  $n \times n$  matrix. Then,  $\det 2(A) = \det_{\psi_2}(A) + (n - 1) \det A$ .*

With this result, we get the following corollary of Theorem 3.5.

**Corollary 3.12.** *Let  $D_G$  be the product distance matrix of a connected graph  $G$  with blocks  $B_1, B_2, \dots, B_p$ , with the property that  $\det D_{B_i} \neq 0$  for all  $1 \leq i \leq p$ . Then,*

$$\det_{\psi_2}(D_G) = \sum_{i=1}^p \left[ \det 2(D_{B_i}) \prod_{j \neq i} \det D_{B_j} \right] - (n - 1) \prod_{i=1}^p \det D_{B_i}.$$

*In particular,  $\det_{\psi_2}(D_G)$  is independent of the tree-like manner of connection of  $G$ 's blocks.*

#### 4. Corollaries

In this section, we derive some corollaries for the exponential distance matrix of a tree  $T$  and the  $q$ -analogue of  $T$ 's laplacian. We then give a  $q$ -analogue again for trees of Schur's dominance theorem. Lastly, we find  $\det 2(K)$  where  $K$  is the matrix appearing in Theorem 3.1.

**4.1. The  $q$ -analogue of  $T$ 's laplacian.** Let  $G = T$  be a tree with exponential distance matrix  $\text{ED}_T$ . Let  $A$  be the adjacency matrix of  $T$  and  $D$  be a diagonal matrix with  $d_{v,v} = \deg(v)$  where  $\deg(v)$  is the degree of vertex  $v$  in  $T$ . Define the  $q$ -analogue of  $T$ 's laplacian as  $\mathcal{L}_q = I - qA + q^2(D - I)$  where  $q$  is an indeterminate. It is easy to see that when  $q = 1$ ,  $\mathcal{L}_q = L$ , where  $L$  is the laplacian matrix of  $T$ .

Bapat, Lal and Pati (see [1, Proposition 3.3]) showed the following surprising inverse result for trees. Their result can alternatively be proved using Theorem 3.1.

**Theorem 4.1.** *Let  $T$  be a tree with exponential distance matrix  $\text{ED}_T$ . Then  $\text{ED}_T^{-1} = \frac{1}{1 - q^2} \mathcal{L}_q$ .*

Theorem 4.1 gives us the following corollary.

**Corollary 4.2.** *Let  $T$  be a tree on  $n \geq 2$  vertices and let  $\text{ED}_T$  be its exponential distance matrix. Then,  $\det 2(\text{ED}_T) = (n - 1)(1 - q^2)^{n-2}(1 + q^2)$ .*

**Proof 4.3.** *There are several ways to prove this. We give two of them. From Theorem 4.1, we see that  $\det \text{ED}_T(i) = (1 - q^2)^{n-2}[1 + q^2(\deg(i) - 1)]$ . It is known that  $\det \text{ED}_T = (1 - q^2)^{n-1}$ . Plugging both of these in Theorem 1.1, we get  $\det 2(\text{ED}_T) = (n - 1)(1 - q^2)^{n-2}(1 + q^2)$ .*

*Alternatively, we use Theorem 3.5. Each block of  $T$  is  $K_2$ , the complete graph on 2 vertices and there are  $n - 1$  such blocks. Thus, for all  $1 \leq i < n$ ,  $D_{B_i} = \begin{pmatrix} 1 & q \\ q & 1 \end{pmatrix}$ . It is simple to see that  $\det D_{B_i} = 1 - q^2$  and that  $\det 2(D_{B_i}) = 1 + q^2$ . The proof is complete by applying Theorem 3.5.*

The second immanant  $\det 2(\mathcal{L}_q)$  can also be found using the above result. We recall the following striking theorem of Merris (see Merris and Watkins [9, Page 239]).

**Theorem 4.4** (Merris). *Let  $A$  be an invertible matrix. Then,  $\det A \cdot \det 2(A^{-1}) = \det A^{-1} \cdot \det 2(A)$ .*

Let  $T$  be a tree with  $\text{ED}_T$  as its exponential distance matrix and let  $\mathcal{L}_q$  be the  $q$ -analogue of its laplacian. Below, we present a  $q$ -analogue of Corollary 2.2.

**Corollary 4.5.** *If  $T$  is a tree on  $n$  vertices and  $\mathcal{L}_q$  is the  $q$ -analogue of its laplacian, then,  $\det 2(\mathcal{L}_q) = (n - 1)(q^2 + 1)$ . Hence,  $\det 2(\mathcal{L}_q)$  only depends on  $n$ , and is independent of the structure of the tree  $T$ .*

**Proof 4.6.** *It is known (see [1]), that  $\det \text{ED}_T = (1 - q^2)^{n-1}$ ,  $\det \mathcal{L}_q = (1 - q^2)$ . Using Theorems 4.4, 4.1 and combining this with Corollary 4.2 yields  $\det 2(\mathcal{L}_q) = (n - 1)(q^2 + 1)$ . Alternatively, one can just use Theorems 4.1 and 1.1.*

We recall the following notation used in  $q$ -series theory. Let  $q$  be an indeterminate and for a positive integer  $i$ , let  $[i]_q = 1 + q + \dots + q^{i-1}$  with  $[0]_q = 0$ . Then, the above corollary can be alternatively

written as  $\det 2(\mathcal{L}_q) = (n - 1)[2]_{q^2}$  for trees. Clearly, setting  $q = 1$  gives us  $[2]_{q^2} = 2$  and  $\mathcal{L}_q = L$ . In this case, we recover Corollary 2.2.

**4.2. Schur’s Dominance Theorem.** Schur [11] showed the following “Dominance Theorem” for positive semidefinite matrices  $A = (a_{i,j})_{1 \leq i,j \leq n}$ . Recall that if  $\lambda$  is an irreducible representation of  $\mathfrak{S}_n$  over the complex numbers  $\mathbb{C}$  with character  $\chi_\lambda$ , then,  $\det_{\chi_\lambda}(A) = \sum_{\sigma \in \mathfrak{S}_n} \chi_\lambda(\sigma) \prod_{i=1}^n a_{i,\sigma_i}$ . Also recall that  $\chi_\lambda(\text{id})$  is the degree of the representation  $\lambda$ , where  $\text{id}$  is the identity permutation in  $\mathfrak{S}_n$ . With these, Schur’s result can be stated as follows.

**Theorem 4.7.** *Let  $A$  be an  $n \times n$  positive semidefinite matrix and let  $\lambda$  be an irreducible representation of  $\mathfrak{S}_n$  with character  $\chi_\lambda$ . Then,  $\det_{\chi_\lambda}(A) \geq \chi_\lambda(\text{id}) \det A$ .*

It is well known that the laplacian matrix  $L$  of any graph is positive semidefinite (see [15]). Similarly, the  $n \times n$  matrix  $J_n$  with all entries being 1 is also positive semidefinite. If  $\lambda_2$  is the irreducible representation of  $\mathfrak{S}_n$  indexed by the partition  $2, 1^{n-2}$ , then by the Hook-length formula, it follows that  $\chi_2(\text{id}) = n - 1$  where  $\chi_2$  denotes  $\chi_{\lambda_2}$ , see Sagan [10]. Thus, if  $T$  is a tree on  $n$  vertices, then it follows that  $\det 2(L) \geq (n - 1) \det L \geq 0$ . As seen earlier for all graphs, when  $q = 1$ ,  $\mathcal{L}_q = L$ . Similarly for trees  $T$  on  $n$  vertices, when  $q = 1$ , then  $\text{ED}_T = J_n$ . Thus, when  $q = 1$ , we have  $\det 2(\mathcal{L}_q) \geq 0$  for all graphs (and hence for trees) and for trees  $T$ , we have  $\det 2(\text{ED}_T) \geq 0$ . We show that for a tree, both  $\text{ED}_T$  and  $\mathcal{L}_q$  satisfy Theorem 4.7 for all  $q \in \mathbb{R}$ .

**Corollary 4.8.** *Let  $T$  be a tree on  $n$  vertices. Let  $\mathcal{L}_q$  be the  $q$ -analogue of its laplacian and let  $\text{ED}_T$  be its exponential distance matrix. Then, for all  $q \in \mathbb{R}$ ,  $\det 2(\mathcal{L}_q) \geq \chi_2(\text{id}) \det(\mathcal{L}_q)$  and for  $-1 \leq q \leq 1$ ,  $\det 2(\text{ED}_T) \geq \chi_2(\text{id}) \det(\text{ED}_T)$ .*

**Proof 4.9.** *From the proof of Corollary 4.5, we have  $\det 2(\mathcal{L}_q) = (n - 1)(1 + q^2)$ ,  $\det \mathcal{L}_q = (1 - q^2)$  and  $\chi_2(\text{id}) = (n - 1)$ . Similarly, Corollary 4.2 gives us  $\det 2(\text{ED}_T) = (n - 1)(1 - q^2)^{n-2}(1 + q^2)$  and  $\det \text{ED}_T = (1 - q^2)^{n-1}$ . Plugging in these values completes the proof.*

**4.3.  $\det 2(\mathcal{D}_T^{-1})$ .** In this subsection, we find  $\det 2(\mathcal{D}_T^{-1})$ , where  $\mathcal{D}_T$  is the non-commutative analogue of the distance matrix of  $T$ . Using Theorem 4.4 and Theorem 1.3, we get the following corollary. Recall the following notation from Theorem 3.8. If  $W_i$  is the  $s \times s$  “weight matrix” on edge  $e_i$ , then  $L_i = \begin{pmatrix} I & W_i \\ W_i & I \end{pmatrix}$ . It is simple to see that  $\det L_i = \det(I - W_i^2)$ .

**Corollary 4.10.** *For a tree  $T$ , let  $\mathcal{D}_T$  be the non-commutative analogue of its distance matrix and let  $K = \mathcal{D}_T^{-1}$  be its inverse. Then,*

$$(4.1) \quad \det 2(K) = \frac{\sum_{i=1}^{n-1} \frac{\det 2(L_i)}{\det L_i} - (n - 2)(s - 1)}{\det \mathcal{D}_T}$$

**Proof 4.11.** From Theorem 4.4, we get  $\det 2(A^{-1}) = \frac{\det 2(A)}{(\det A)^2}$ . Theorem 1.3 gives a product rule that  $\det \mathcal{D}_T = \prod_{i=1}^{n-1} \det L_i$  and Theorem 3.8 gives us

$$\det 2(\mathcal{D}_T) = \left( \sum_{i=1}^{n-1} \det 2(L_i) \prod_{j \neq i} \det L_j \right) - (n-2)(s-1) \det \mathcal{D}_T.$$

Combining these completes the proof.

### Acknowledgements

Some Theorems in this work were as conjectures, tested using the computer package ‘‘Sage’’. We thank the authors of ‘‘Sage’’ for generously releasing their software as an open-source package. The first author acknowledges support from the JC Bose Fellowship, Department of Science and Technology, Government of India. The second author acknowledges support from project grant P07 IR052, given by IIT Bombay.

We thank the referee for several suggestions which has improved the readability of this work.

### REFERENCES

- [1] R. B. Bapat, A. K. Lal and S. Pati, A  $q$ -analogue of the distance matrix of a tree, *Linear Algebra Appl.*, **416** (2006) 799–814.
- [2] R. B. Bapat and S. Sivasubramanian, Product Distance Matrix of a Graph and Squared Distance Matrix of a Tree, *Appl. Anal. Discrete Math.*, **7** (2013) 285–301.
- [3] R. B. Bapat and S. Sivasubramanian, Product distance matrix of a tree with matrix weights, *Linear Algebra Appl.*, **468** (2015) 145–153.
- [4] R. L. Graham, A. J. Hoffman and H. Hosoya, On the distance matrix of a directed graph, *J. Graph Theory*, **1** (1977) 85–88.
- [5] R. L. Graham and H. O. Pollak, On the addressing problem for loop switching, *Bell System Tech. J.*, **50** (1971) 2495–2519.
- [6] R. Grone and R. Merris, An Algorithm for the Second Immanant, *Maths. Comp.*, **43** no. 168 (1984) 589–591.
- [7] D. E. Littlewood, *The theory of group characters and matrix representations of groups*, Reprint of the second (1950) edition, Providence, RI, 2006.
- [8] R. Merris, The second immanantal polynomial and the centroid of a graph, *SIAM J. Algebraic Discrete Methods*, **7** (1986) 484–503.
- [9] R. Merris and W. Watkins, Inequalities and identities for generalized matrix functions, *Linear Algebra Appl.*, **64** (1985) 223–242.
- [10] B. E. Sagan, *The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions*, Second edition. Graduate Texts in Mathematics, **203**, Springer-Verlag, New York, 2001.
- [11] I. Schur, Über endliche Gruppen und Hermitesche Formen, *Math. Z.*, **1** (1918) 184–207.
- [12] S. Sivasubramanian, A  $q$ -analogue of Graham, Hoffman and Hosoya’s theorem, *Electron. J. Combin.*, **17** (2010) Note 21 9 pp.
- [13] R. P. Stanley, *Enumerative Combinatorics*, **2**, Cambridge University Press, Cambridge, 1999.
- [14] J. R. Stembridge, Some Conjectures For Immanants, *Canad. J. Math.*, **44** (1992) 1079–1099.

[15] D. B. West, *Introduction to Graph Theory, 2nd Edition*, Prentice Hall, Inc., Upper Saddle River, NJ, 1996.

**R. B. Bapat**

Stat-Math Unit, Indian Statistical Institute, New Delhi, India

Email: [rbb@isid.ac.in](mailto:rbb@isid.ac.in)

**Sivaramakrishnan Sivasubramanian**

Department of Mathematics, Indian Institute of Technology, Bombay, Mumbai, India

Email: [krishnan@math.iitb.ac.in](mailto:krishnan@math.iitb.ac.in)